

HOMOLOGY OF THE MODULI SPACES AND MAPPING CLASS GROUPS OF FRAMED, r -SPIN AND PIN SURFACES

OSCAR RANDAL-WILLIAMS

ABSTRACT. We give definitions of moduli spaces of framed, r -Spin and $\text{Pin}^\pm$ surfaces. We apply earlier work of the author to show that each of these moduli spaces exhibits homological stability, and we identify the stable integral homology with that of certain infinite loop spaces in each case. We further show that these moduli spaces each have path components which are Eilenberg–MacLane spaces for the framed, r -Spin and $\text{Pin}^\pm$ mapping class groups respectively, and hence we also identify the stable group homology of these groups.

In particular: the stable framed mapping class group has trivial rational homology, and its abelianisation is $\mathbb{Z}/24$; the rational homology of the stable $\text{Pin}^\pm$ mapping class groups coincides with that of the non-orientable mapping class group, and their abelianisations are $\mathbb{Z}/2$ for Pin^+ and $(\mathbb{Z}/2)^3$ for Pin^- .

1. INTRODUCTION AND STATEMENT OF RESULTS

Recent advances in the theory of moduli spaces of complex curves [14, 10] concern the stable topology of these spaces, that is, the topology of moduli spaces of curves of high genus. This theory has been built upon by several workers to—amongst other things—deal with tangential structures other than orientations. Thus much is known about the homology of moduli spaces of unoriented surfaces [21], moduli spaces of Spin surfaces [11, 1, 7], and moduli spaces of oriented surfaces with maps to a simply-connected background space [2, 3].

The author has recently given [18] a general theory of homological stability for moduli spaces of surfaces with θ -structure (which we define below). This recovers the above examples, but also allows one to effectively study many moduli spaces that are new to the literature. In this paper we study three examples of these: moduli spaces of framed surfaces, moduli spaces of r -Spin surfaces, and moduli spaces of $\text{Pin}^\pm$ surfaces.

Let us give a precise definition of the moduli spaces we have in mind. Write F for a smooth surface, possibly with boundary. Let $\theta : \mathbf{X} \rightarrow BO(2)$ be a Serre fibration classifying the bundle $\theta^*\gamma_2 \rightarrow \mathbf{X}$, and let $\text{Bun}(TF, \theta^*\gamma_2)$ denote the space of bundle maps $TF \rightarrow \theta^*\gamma_2$, i.e. fibrewise linear isomorphisms. Given a bundle map $\delta : TF|_{\partial F} \rightarrow \theta^*\gamma_2$, let $\text{Bun}_\partial(TF, \theta^*\gamma_2; \delta)$ denote the space of bundle maps $TF \rightarrow \theta^*\gamma_2$ that restrict to δ on the boundary. Let $\text{Diff}_\partial(F)$ denote the group of diffeomorphisms of F which restrict to the identity diffeomorphism on some neighbourhood of the boundary, and equip it with the C^∞ topology.

Date: February 7, 2011.

2010 Mathematics Subject Classification. 55R40, 57R15, 57R50, 57M07.

Key words and phrases. Mapping class groups, Moduli spaces, Surface bundles.

The author has been supported by an EPSRC Studentship, DTA grant number EP/P502667/1, ERC Advanced Grant No. 228082, and the Danish National Research Foundation through the Centre for Symmetry and Deformation.

Definition 1.1. The *moduli space of surfaces with θ -structure of topological type F and boundary condition δ* is the homotopy quotient

$$\mathcal{M}^\theta(F; \delta) := \text{Bun}_\partial(TF, \theta^*\gamma_2; \delta) // \text{Diff}_\partial(F).$$

If we define $\mathcal{E}^\theta(F; \delta) := \text{Bun}_\partial(TF, \theta^*\gamma_2; \delta) \times F // \text{Diff}_\partial(F)$, where the group acts diagonally, the projection map

$$\mathcal{E}^\theta(F; \delta) \longrightarrow \mathcal{M}^\theta(F; \delta)$$

is a smooth F -bundle equipped with a bundle map $T^v\mathcal{E}^\theta(F; \delta) \rightarrow \theta^*\gamma_2$ from the vertical tangent bundle satisfying appropriate boundary conditions, and is universal with this property. This construction is discussed in more depth in [8, 9, 18].

Definition 1.2. Given a θ -structure δ on ∂F , and a point $\xi \in \mathcal{M}^\theta(F; \delta)$, we define the *θ mapping class group* (at ξ) to be the fundamental group

$$\Gamma^\theta(F; \xi) := \pi_1(\mathcal{M}^\theta(F; \delta), \xi).$$

If we do not wish to introduce notation for a boundary condition, we may write $\mathcal{M}^\theta(F)$ to denote $\mathcal{M}^\theta(F; \delta)$ with an unspecified δ .

These moduli spaces have certain *stabilisation maps* between them. Let F be a surface with boundary condition $\delta : TF|_{\partial F} \rightarrow \theta^*\gamma_2$ and F' be a surface with θ -structure. Given an identification of manifolds with θ -structure from a collection of boundary components of F' to a collection of boundary components of F , there is a map

$$\mathcal{M}^\theta(F; \delta) \longrightarrow \mathcal{M}^\theta(F \cup F'; \delta')$$

obtained by gluing F' to F along the identified boundaries.

If we write $\Sigma_{g,b}$ for the orientable surface of genus g with b boundary components, these stabilisation maps for orientable surfaces are generated by certain elementary stabilisation maps

$$\begin{aligned} \alpha(g) : \mathcal{M}^\theta(\Sigma_{g,b}; \delta) &\longrightarrow \mathcal{M}^\theta(\Sigma_{g+1,b-1}; \delta') \\ \beta(g) : \mathcal{M}^\theta(\Sigma_{g,b}; \delta) &\longrightarrow \mathcal{M}^\theta(\Sigma_{g,b+1}; \delta') \\ \gamma(g) : \mathcal{M}^\theta(\Sigma_{g,b}; \delta) &\longrightarrow \mathcal{M}^\theta(\Sigma_{g,b-1}; \delta') \end{aligned}$$

given by gluing on a pair of pants along the legs, a pair of pants along the waist, and a disc, respectively. If we write $S_{n,b}$ for the non-orientable surface of genus n with b boundary components, in addition to the analogues of the above maps there are also stabilisation maps

$$\mu(n) : \mathcal{M}^\theta(S_{n,b}; \delta) \longrightarrow \mathcal{M}^\theta(S_{n+1,b}; \delta')$$

given by gluing on a projective plane with two discs removed.

Theorems about the stable topology of the moduli spaces $\mathcal{M}^\theta(F)$ typically take the form of stating that a comparison map to a certain infinite loop space is a homology equivalence in some range of degrees.

Definition 1.3. The *Madsen–Tillmann spectrum* of θ , denoted $\mathbf{MT}\theta$, is the Thom spectrum of the virtual bundle $-\theta^*\gamma_2 \rightarrow \mathbf{X}$. We denote by $\Omega^\infty \mathbf{MT}\theta$ the associated infinite loop space.

For closed surfaces F there is a natural comparison map

$$\alpha_F : \mathcal{M}^\theta(F) \longrightarrow \Omega^\infty \mathbf{MT}\theta$$

defined using Pontrjagin–Thom theory, and many characteristic classes of θ -surface bundles exist universally (i.e. independent of the topological type of F) on $\Omega^\infty \mathbf{MT}\theta$.

1.1. Moduli spaces of framed surfaces. Let us take the tangential structure $\theta : EO(2) \rightarrow BO(2)$, and write $\mathcal{M}^{\text{fr}}(\Sigma_{g,b}; \delta)$ for the moduli space of framed surfaces with underlying surface $\Sigma_{g,b}$. Write $\Gamma^{\text{fr}}(\Sigma_{g,b}; \xi)$ for the framed mapping class group. We show that the path component of $\mathcal{M}^{\text{fr}}(\Sigma_{g,b}; \delta)$ containing ξ is homotopy equivalent to $B\Gamma^{\text{fr}}(\Sigma_{g,b}; \xi)$, so homological questions about the framed mapping class group are equivalent to homological questions about the moduli space of framed surfaces.

Our main theorem concerning the moduli spaces of framed surfaces is that they exhibit homological stability: the homology groups $H_*(\mathcal{M}^{\text{fr}}(\Sigma_{g,b}; \delta); \mathbb{Z})$ are independent of g , b and δ as long as $5* \leq 2g - 7$. Furthermore, the stable homology coincides with the homology of the space $\Omega^\infty \mathbf{MTEO}(2) = \Omega^\infty \mathbf{S}^{-2} \simeq \Omega^2 Q_0 S^0$, the double loop space of the free infinite loop space on a point. In particular, it follows that the abelianisation of the group $\Gamma^{\text{fr}}(\Sigma_{g,b}; \xi)$ is $\mathbb{Z}/24$ as long as $g \geq 6$, and that the rational group homology of $\Gamma^{\text{fr}}(\Sigma_{g,b}; \xi)$ is trivial in the stable range.

1.2. Moduli spaces of r -Spin surfaces. Recall that $\text{Spin}^r(2) = SO(2)$, but its standard 1-dimensional (complex) representation is the r -th tensor power of the standard representation of $SO(2)$.

Fixing an r , we take the tangential structure $\theta : B\text{Spin}^r(2) \rightarrow BO(2)$, and write $\mathcal{M}^{\text{Spin}^r}(\Sigma_{g,b}; \delta)$ for the moduli space of r -Spin surfaces with underlying surface $\Sigma_{g,b}$. Write $\Gamma^{\text{Spin}^r}(\Sigma_{g,b}; \xi)$ for the r -Spin mapping class group. We show that the component of $\mathcal{M}^{\text{Spin}^r}(\Sigma_{g,b}; \delta)$ containing ξ is homotopy equivalent to $B\Gamma^{\text{Spin}^r}(\Sigma_{g,b}; \xi)$, so homological questions about the r -Spin mapping class group are equivalent to homological questions about the moduli space of r -Spin surfaces.

Our main theorem concerning the moduli spaces of r -Spin surfaces is that they exhibit homological stability: the homology groups $H_*(\mathcal{M}^{\text{Spin}^r}(\Sigma_{g,b}; \delta); \mathbb{Z})$ are independent of g , b and δ in degrees $5* \leq 2g - 7$.

In the case $r = 2$, our 2-Spin mapping class groups coincide with the extended Spin mapping class groups of Masbaum [15]. Galatius has shown [7] that the stable homology of these groups coincides with the homology of the infinite loop space $\Omega_0^\infty \mathbf{MTSpin}(2)$. We will show that a similar description of the stable homology is possible for all r , and give computational applications of this result, in [19].

1.3. Moduli spaces of $\text{Pin}^\pm$ surfaces. Recall that there are two generalisations of the covering group $\text{Spin}(2) \rightarrow SO(2)$ to a covering group of $O(2)$, called $\text{Pin}^+(2)$ and $\text{Pin}^-(2)$, which we recall in §4. These give Serre fibrations $\theta : B\text{Pin}^\pm(2) \rightarrow BO(2)$ with corresponding moduli spaces $\mathcal{M}^{\text{Pin}^+}(S_{n,b}; \delta)$ and $\mathcal{M}^{\text{Pin}^-}(S_{n,b}; \delta)$, and mapping class groups $\Gamma^{\text{Pin}^+}(S_{n,b}; \xi)$ and $\Gamma^{\text{Pin}^-}(S_{n,b}; \xi)$. We show that the component of $\mathcal{M}^{\text{Pin}^\pm}(S_{n,b}; \delta)$ containing a point ξ is homotopy equivalent to $B\Gamma^{\text{Pin}^\pm}(S_{n,b}; \xi)$, so homological questions about the $\text{Pin}^\pm$ mapping class group are equivalent to homological questions about the $\text{Pin}^\pm$ moduli space.

Our main theorems concerning these moduli spaces is that they exhibit homological stability (for non-orientable surfaces), and we give precise stability ranges in §4.6. Furthermore, the stable homology coincides with that of the infinite loop spaces $\Omega^\infty \mathbf{MTPin}^+(2)$ and $\Omega^\infty \mathbf{MTPin}^-(2)$ respectively. In particular, we are able to calculate that the abelianisation of the group $\Gamma^{\text{Pin}^+}(S_{n,b}; \delta)$ is $\mathbb{Z}/2$ as long as $n \geq 9$, and that of $\Gamma^{\text{Pin}^-}(S_{n,b}; \delta)$ is $(\mathbb{Z}/2)^3$ as long as $n \geq 11$. In §5.1 we also study the divisibility of certain characteristic classes ζ_i which were defined by Wahl in the integral cohomology of the moduli spaces of unoriented surfaces, when they are pulled back to the moduli spaces $\mathcal{M}^{\text{Pin}^\pm}(F)$.

1.4. Guide. We prove the homological stability theorems for these tangential structures by applying the general stability theorems of [18]. In order to verify the hypotheses of these theorems it is necessary to obtain a good understanding of the sets of path components $\pi_0(\mathcal{M}^\theta(F; \delta))$ and the effect of stabilisation maps between these sets. The bulk of the paper is dedicated to this problem for the tangential structures in question, and can be understood without reference to [18]. In the proofs of the homological stability theorems we refer to concepts defined in [18, §5–9], and we will not give these definitions again.

The main qualitative theorem of [18] is that if the sets $\pi_0(\mathcal{M}^\theta(F; \delta))$ stabilise then the homology of these spaces stabilises. Much of the trouble we go to in this paper is to find the best possible stability range that the methods of that paper can provide.

2. MODULI SPACES OF FRAMED AND r -SPIN SURFACES

In the introduction we explained how a map $\theta : \mathbf{X} \rightarrow BO(2)$ produces a moduli space of surfaces with θ -structure $\mathcal{M}^\theta(F; \delta)$ for each surface F and boundary condition δ .

Definition 2.1. To define $\mathcal{M}^{\text{fr}}(F; \delta)$, the *moduli space of framed surfaces* of topological type F and boundary condition δ , we take the map $\theta_0 : EO(2) \rightarrow BO(2)$ which classifies γ_2^0 , the trivialised bundle.

To define $\mathcal{M}^{\text{Spin}^r}(F; \delta)$, the *moduli space of r -Spin surfaces* of topological type F and boundary condition δ , we take the map $\theta_r : B\text{Spin}^r(2) \rightarrow BO(2)$, where $B\text{Spin}^r(2) := BU(1)$ and the map to $BO(2)$ classifies γ_2^r , the r -th power of the tautological complex line bundle.

The maps defining both of these tangential structures naturally factor through the map $\theta_+ : BSO(2) \rightarrow BO(2)$ classifying γ_2^+ , the universal oriented rank 2 vector bundle, and so they have forgetful maps which just remember the orientation. If $\ell : TF \rightarrow \gamma_2^r$ is a θ_r -structure on F , we denote by $\ell^+ : TF \rightarrow \gamma_2^+$ the underlying orientation obtained from the map $\gamma_2^r \rightarrow \gamma_2^+$. This defines a map

$$\text{Bun}_\partial(TF, \gamma_2^r; \delta) \longrightarrow \text{Bun}_\partial(TF, \gamma_2^+; \delta^+)$$

whose fibres are either empty (over a θ_+ -structure which does not admit a refinement to a θ_r -structure), or else are homotopy equivalent to $\text{map}_*(F/\partial F, B\mathbb{Z}/r)$, where if $\partial F = \emptyset$ we interpret $F/\partial F$ as F with a disjoint basepoint adjoined. Hence there are fibrations

$$(2.1) \quad \text{map}_*(F/\partial F, SO(2)) \longrightarrow \mathcal{M}^{\text{fr}}(F; \delta) \longrightarrow \mathcal{M}^+(F; \delta^+)$$

$$(2.2) \quad \text{map}_*(F/\partial F, B\mathbb{Z}/r) \longrightarrow \mathcal{M}^{\text{Spin}^r}(F; \delta) \longrightarrow \mathcal{M}^+(F; \delta^+)$$

whenever the total space is non-empty. Note that if F has no boundary, $\mathcal{M}^{\text{fr}}(F; \delta)$ is non-empty if and only if F is diffeomorphic to a torus. Thus when discussing framed surfaces we will always suppose that they have boundary. Furthermore, from now on we will assume that all surfaces are orientable.

2.1. Naturality properties. If r' divides r , there is a map of moduli spaces

$$\mathcal{M}^{\text{Spin}^r}(F; \delta_r) \longrightarrow \mathcal{M}^{\text{Spin}^{r'}}(F; \delta_{r'}),$$

where $\delta_{r'}$ is the induced r' -Spin structure from the r -Spin structure δ_r . Furthermore, there are maps to all of these moduli spaces from $\mathcal{M}^{\text{fr}}(F; \delta)$.

2.2. Mapping class groups. In the introduction we defined the θ mapping class group of a θ -surface $\xi \in \mathcal{M}^\theta(F; \delta)$ to be the fundamental group based at this point.

In the case of framings—as ∂F is assumed to be non-empty—in the fibration (2.1) the space $\text{map}_*(F/\partial F, SO(2))$ is homotopy-discrete and so we have an exact sequence of groups and pointed sets

$$0 \longrightarrow \pi_1(\mathcal{M}^{\text{fr}}(F; \delta), \xi) \longrightarrow \Gamma^+(F) \xrightarrow{\varphi} H^1(F, \partial F; \mathbb{Z}) \longrightarrow \pi_0(\mathcal{M}^{\text{fr}}(F; \delta)) \longrightarrow *.$$

Here $\Gamma^+(F)$ denotes the usual mapping class group of the oriented surface F , and the map φ coincides with the crossed homomorphism obtained by Trapp [20] which gives an extended symplectic representation of the oriented mapping class group, though we will not pursue this connection. The above sequence identifies the framed mapping class group $\Gamma^{\text{fr}}(F; \xi)$ as the subgroup of the oriented mapping class group $\Gamma^+(F)$ consisting of those (isotopy classes of) diffeomorphisms which fix the isotopy class of framings $[\xi]$. Furthermore, it implies that $\mathcal{M}^{\text{fr}}(F; \delta)$ is a disjoint union of $K(\pi, 1)$'s, i.e. has the homotopy type of a groupoid.

In the case of r -Spin structures, if we suppose that ∂F is non-empty, the space $\text{map}_*(F/\partial F, B\mathbb{Z}/r)$ is homotopy-discrete so the fibre sequence (2.2) gives the exact sequence of groups and pointed sets

$$0 \longrightarrow \pi_1(\mathcal{M}^{\text{Spin}^r}(F; \delta), \xi) \longrightarrow \Gamma^+(F) \xrightarrow{\varphi} H^1(F, \partial F; \mathbb{Z}/r) \longrightarrow \dots$$

This identifies the r -Spin mapping class group $\Gamma^{\text{Spin}^r}(F; \xi)$ as the subgroup of the oriented mapping class group $\Gamma^+(F)$ consisting of those (isotopy classes of) diffeomorphisms which fix the isotopy class of r -Spin structures $[\xi]$.

If F does not have boundary, there is a sequence

$$0 \longrightarrow \mathbb{Z}/r \longrightarrow \pi_1(\mathcal{M}^{\text{Spin}^r}(F; \delta), \xi) \longrightarrow \Gamma^+(F) \xrightarrow{\varphi} H^1(F; \mathbb{Z}/r) \longrightarrow \dots$$

which identifies the r -Spin mapping class group with an extension by $\mathbb{Z}/r$ of the subgroup of $\Gamma^+(F)$ of mapping classes that preserve a r -Spin structure up to isomorphism. In either case, $\mathcal{M}^{\text{Spin}^r}(F; \delta)$ is a disjoint union of $K(\pi, 1)$'s, i.e. has the homotopy type of a groupoid.

2.3. The set of θ_r -structures. In order to apply the results of [18], we must calculate the set of path components $\pi_0(\mathcal{M}^{\theta_r}(F; \delta))$, that is, calculate the set of isotopy classes of θ_r -structures a surface admits up to diffeomorphism of the underlying surface. This coincides with the quotient set of $\pi_0 \text{Bun}_\partial(TF, \gamma_2^{\text{fr}}; \delta)$ or $\pi_0 \text{Bun}_\partial(TF, \gamma_2^r; \delta)$ by the action of $\Gamma^+(F)$.


Let us define

$$\theta_r(F; \delta) := \begin{cases} \pi_0 \text{Bun}_\partial(TF, \gamma_2^{\text{fr}}; \delta) & \text{if } r = 0 \\ \pi_0 \text{Bun}_\partial(TF, \gamma_2^r; \delta) & \text{if } r > 0. \end{cases}$$

Definition 2.2. Choose once and for all a θ_r -structure on $\mathbb{R}^2$, that is, a linear map $\mathbb{R}^2 \rightarrow \gamma_2^r$. If $V \rightarrow B$ is a framed rank two vector bundle, the *standard θ_r -structure* is the fibrewise linear isomorphism $V \rightarrow \mathbb{R}^2 \rightarrow \gamma_2^r$.

On the surface $\Sigma_{g,b+1}$ having θ_r -structure ξ with boundary condition δ , choose a marked point on each boundary component, and equip each marked point with the framing coming from the orientation and the inwards pointing normal vector. After perhaps changing δ to an isomorphic boundary condition, we may suppose that the θ_r -structure is standard at each marked point.

Let $\{a_i, b_i, t_i, \partial_i, r_i\}$ be the collection of simple closed curves and simple arcs in $\Sigma_{g,b+1}$ as shown in Figure 1, where the endpoints of the arcs lie at the standard marked points on each boundary component. Note that $a_i \cap b_j = \delta_{ij}$ and $a_i \cap a_j = b_i \cap b_j = 0$. These curves have canonical lifts $\tilde{a}_i, \tilde{b}_i, \tilde{t}_i, \tilde{\partial}_i, \tilde{r}_i$ to curves or arcs on $S\Sigma_{g,b+1}$, given by assigning them their unit forwards tangent vector at each point.

FIGURE 1. Curves and arcs on $\Sigma_{g,b+1}$.

We remark that these lifts are *not* homology invariant: although t_1 is homologous to $a_2 - a_1$, $\tilde{t}_1$ is homologous to $\tilde{a}_2 - \tilde{a}_1 + z$. In general, when lifting homologous elements from $\Sigma_{g,b+1}$ to $\mathbb{S}\Sigma_{g,b+1}$, there is a correction term given by the Euler characteristic of a homology chain times z .

Definition 2.3. Given a θ_r -structure ξ on $\Sigma_{g,b+1}$ we define a $\mathbb{Z}/r$ -valued function q_ξ on the set of simple closed curves (or simple arcs between the standard marked points on each boundary component) by assigning to each curve or arc x the value $q_\xi(x)$ determined as follows: $TF|_x$ has a θ_r -structure, but the forwards vector field along x splits off a trivial 1-dimensional sub-bundle, so reduces the structure group to $\text{Spin}^r(1) = \mathbb{Z}/r$. If x is a simple closed curve, the monodromy gives an element of $\mathbb{Z}/r$. If x is an arc, the forward / left framing along the arc agrees with the standard framing at the start of the arc, but not at the end: here it differs by a half rotation. However, the orientation of the surface gives a canonical choice of half rotation, which makes the θ_r -structure on the arc be standard near its ends, and hence give an element of $\mathbb{Z}/r$. In both cases we denote the element of $\mathbb{Z}/r$ obtained by $\text{Mon}(x)$.

We define $q_\xi(x)$ to be $\text{Mon}(x) - 1 \in \mathbb{Z}/r$.

The reader may be at a loss as to why we subtract 1 to what is already a perfectly good invariant: it is a normalisation, and ensures that if a simple closed curve x bounds a disc, then $q_\xi(x) = 0$. We may now define a function

$$p : \theta_r(\Sigma_{g,b+1}; \delta) \longrightarrow (\mathbb{Z}/r)^{2g+b}$$

given by $p(\xi) := (q_\xi(a_1), q_\xi(b_1), \dots, q_\xi(a_g), q_\xi(b_g), q_\xi(r_1), \dots, q_\xi(r_b))$.

Proposition 2.4. *If $\theta_r(\Sigma_{g,b+1}; \delta)$ is non-empty, the map p is a bijection.*

Proof. Note that once a θ_r -structure is determined over the curves a_i , b_i and r_j , it remains to give a θ_r -structure on a disc satisfying a certain boundary condition (which up to isomorphism depends only on δ , and not on the values of ξ on the curves). This is possible (and if so, in a unique way) if and only if $\theta_r(\Sigma_{g,b+1}; \delta)$ is non-empty, by obstruction theory for the map θ_r . $\square$

Using this proposition, we may study the action of the mapping class group $\Gamma^+(\Sigma_{g,b+1})$ on the set $\theta_r(\Sigma_{g,b+1}; \delta)$ by studying its action in the “coordinate system” given by p .

Remark 2.5. We should mention to what extent the function p is unique. We have only defined it for boundary conditions with a choice of standard marked point on each boundary, and an ordering of the boundaries. Varying either of these choices will vary the function.

Lemma 2.6. *A θ_r -structure ξ on F gives an element $[\xi] \in H^1(\mathbb{S}F; \mathbb{Z}/r)$, such that the function q_ξ on simple closed curves or arcs with standard ends is given by lifting it to $\mathbb{S}F$ using the forwards tangent vector, and applying $[\xi]$ and subtracting 1.*

Proof. By a Serre spectral sequence calculation, there is a unique class in $H^1(\mathbb{S}\gamma_2^r; \mathbb{Z}/r)$ which restricts to $1 \in H^1(S^1; \mathbb{Z}/r)$, as the Euler class of γ_2^r is divisible by r . This gives a map $\mathbb{S}\gamma_2^r \rightarrow B\mathbb{Z}/r$ which we see is a homotopy equivalence. A θ_r -structure on F gives a map $\mathbb{S}F \rightarrow \mathbb{S}\gamma_2^r$ taking the standard part of the boundary to the base-point and we define $[\xi]$ to be the pullback of the tautological class via this map. The claimed property is now immediate. $\square$

Let us write τ_a for the (forward) Dehn twist around a simple closed curve a , and recall that the action of a twist on a homology class x is given by the formula $\tau_a(x) = x + \langle a, x \rangle \cdot a$, where $\langle \cdot, \cdot \rangle$ denotes the intersection product.

Lemma 2.7. *If x is a simple closed curve or simple arc on F and $\tilde{x}$ is its canonical lift to $\Sigma\mathbb{S}$ give by the forwards tangent vector, then $\tau_a(\tilde{x}) = \widetilde{\tau_a(x)}$ which is homologous to $\tilde{x} + \langle a, x \rangle \cdot \tilde{a}$. Hence we have the formula*

$$q_{\tau_a^* \xi}(x) = \tau_a^*(q_\xi)(x) = q_\xi(x) + \langle a, x \rangle (q_\xi(a) + 1)$$

for the action of Dehn twists on θ_r -structures.

Proof. The first part may be seen by constructing an explicit homology in $\mathbb{S}F$ between $\widetilde{\tau_a(x)}$ and $\tilde{x} + \langle a, x \rangle \cdot \tilde{a}$ concentrated near the intersection points of a and x , when we represent them by transverse smooth 1-manifolds. The second part now follows as by Lemma 2.6 the function q_ξ is given by evaluating against a cohomology class on $\mathbb{S}F$ —which is linear—and then subtracting 1. $\square$

It is also useful to observe that if a and b are disjoint simple closed curves and $a \# b$ is the simple closed curve obtained by forming the oriented connected sum of a and b , then

$$\widetilde{a \# b} = \tilde{a} + \tilde{b} + z$$

or equivalently for any θ_r -structure ξ ,

$$q_\xi(a \# b) = q_\xi(a) + q_\xi(b).$$

The same holds when one of a and b is a simple arc. More generally, if x is a simple closed curve whose homology class may be written as $[x] = [a] + [b]$ for simple closed curves a, b , then

$$(2.3) \quad q_\xi(x) = q_\xi(a) + q_\xi(b) + \langle a, b \rangle.$$

This follows from the methods of Johnson [12] (in particular his Theorem 1B).

2.4. Diffeomorphism classes of θ_r -structures. Once we have the bijection $p : \theta_r(\Sigma_{g,b+1}; \delta) \longrightarrow (\mathbb{Z}/r)^{2g+b}$, there is a surjective function $A : \theta_2(\Sigma_{g,b+1}; \delta) \rightarrow \mathbb{Z}/2$ given in terms of this bijection by the formula

$$(2.4) \quad \xi \mapsto \sum_{i=1}^g q_\xi(a_i) \cdot q_\xi(b_i) + \sum_{j=1}^b q_\xi(r_j) \cdot \delta_j.$$

For surfaces with zero or one boundaries, the set of 2-Spin structures may be identified with the set of quadratic refinements of the intersection form [12] via $\xi \mapsto q_\xi$. In that case, the second term of this formula vanishes and A is simply the Arf invariant. Hence we call it the *generalised Arf invariant* for 2-Spin surfaces with boundary. We remind the reader that the bijection p was not canonical (it depended on a choice of trivialised marked point on each boundary and an ordering of the boundaries), and hence A considered as a function on $\theta_2(\Sigma_{g,b+1}; \delta)$ is not canonical either.

Proposition 2.8. *The function A is $\Gamma^+(\Sigma_{g,b+1})$ -invariant.*

Proof. Let x be a simple closed curve represented in homology by $\sum_{i=1}^g X_i a_i + Y_i b_i + \sum_{j=0}^b \lambda_j \partial_j$. Then a 2-Spin structure $\xi = (A_1, B_1, \dots, A_g, B_g, R_1, \dots, R_b)$ evaluated on x gives

$$q_\xi(x) = q_\xi \left(\sum_{i=1}^g X_i a_i + Y_i b_i + \sum_{j=0}^b \lambda_j \partial_j \right) = \sum_{i=1}^g X_i A_i + Y_i B_i + X_i Y_i + \sum_{j=0}^b \lambda_j \delta_i$$

using (2.3). The Dehn twist around this curve satisfies

$$\begin{aligned} q_{\tau_x^* \xi}(a_i) &= A_i + Y_i(q_\xi(x) + 1) \\ q_{\tau_x^* \xi}(b_i) &= B_i + X_i(q_\xi(x) + 1) \\ q_{\tau_x^* \xi}(r_j) &= R_j + (\lambda_0 + \lambda_j)(q_\xi(x) + 1) \end{aligned}$$

so if $q_\xi(x) = 1$ then $q_{\tau_x^* \xi} = q_\xi$ and the invariant is trivially preserved. If $q_\xi(x) = 0$ then

$$\sum_{i=1}^g q_{\tau_x^* \xi}(a_i) \cdot q_{\tau_x^* \xi}(b_i) = \sum_{i=1}^g A_i B_i + \left(\sum_{i=1}^g A_i X_i + B_i Y_i + X_i Y_i \right)$$

and

$$\begin{aligned} \sum_{j=1}^r q_{\tau_x^* \xi}(r_i) \cdot \delta_j &= \sum_{j=0}^r R_j \delta_j + \left(\sum_{j=1}^b (\lambda_0 + \lambda_j) \delta_j \right) \\ &= \sum_{j=0}^r R_j \delta_j + \left(\sum_{j=0}^b \lambda_j \delta_j \right) \end{aligned}$$

and $0 = q_\xi(x) = \sum_{j=0}^b \lambda_j \delta_j + \sum_{i=1}^g A_i X_i + B_i Y_i + X_i Y_i$ so $A(\xi) = A(\tau_x^* \xi)$. $\square$

Thus we have produced a $\Gamma^+(\Sigma_{g,b+1})$ -invariant function $A : \theta_2(\Sigma_{g,b+1}) \rightarrow \mathbb{Z}/2$. This induces $\Gamma^+(\Sigma_{g,b+1})$ -invariant functions on all $\theta_{2n}(\Sigma_{g,b+1})$, $n \geq 0$, by composing with the natural map $\theta_{2n}(\Sigma_{g,b+1}; \delta) \rightarrow \theta_2(\Sigma_{g,b+1}; \delta_2)$. The main result of this section is the following theorem, which determines the number of orbits of $\Gamma^+(\Sigma_{g,b+1})$ acting on $\theta_r(\Sigma_{g,b+1}; \delta)$.

Theorem 2.9. *Let $g \geq 2$. If the set $\theta_r(\Sigma_{g,b+1}; \delta)/\Gamma^+(\Sigma_{g,b+1})$ is not empty:*

- (i) *It consists of a single element, if r is odd.*
- (ii) *It consists of two elements distinguished by the invariant A , if r is even.*

In order to prove this theorem we require the following lemma which describes the action of the mapping class group $\Gamma^+(\Sigma_{g,b+1})$ on the set $\theta_r(\Sigma_{g,b+1}; \delta)$, in the coordinate system given by Proposition 2.4.

Lemma 2.10. *Translating the action of $\Gamma^+(\Sigma_{g,1})$ on $\theta_r(\Sigma_{g,1}; \delta)$ to $(\mathbb{Z}/r)^{2g}$, Dehn twists around the cycles a_i and b_i give*

$$\begin{aligned}\tau_{a_i}^{\pm 1} \cdot (A_1, B_1, \dots, A_g, B_g) &= (A_1, B_1, \dots, B_{i-1}, A_i, B_i \pm (A_i + 1), A_{i+1}, \dots, A_g, B_g), \\ \tau_{b_i}^{\pm 1} \cdot (A_1, B_1, \dots, A_g, B_g) &= (A_1, B_1, \dots, B_{i-1}, A_i \mp (B_i + 1), B_i, A_{i+1}, \dots, A_g, B_g).\end{aligned}$$

Dehn twists around the cycles t_i give that $\tau_{t_i} \cdot (A_1, B_1, \dots, A_g, B_g)$ is

$$(A_1, \dots, A_i, B_i + A_i - A_{i+1} - 1, A_{i+1}, B_{i+1} + A_{i+1} - A_i + 1, A_{i+2}, \dots, B_g).$$

The analogous formulae hold for more than one boundary.

Proof. For the first part, we compute $\tau_{a_i}(q_\xi)(b_i) = q_\xi(\tau_{a_i}(b_i)) = q_\xi(b_i) + q_\xi(a_i) + 1$, and so on. For the second part, note that t_i is homologous to $a_{i+1} - a_i$, and that $\tilde{t}_i$ is homologous to $\tilde{a}_{i+1} - \tilde{a}_i + z$. Thus $q_\xi(t_i) = q_\xi(a_{i+1}) - q_\xi(a_i)$,

$$\tau_{t_i}(q_\xi)(b_i) = q_\xi(b_i) - (q_\xi(t_i) + 1) = q_\xi(b_i) + q_\xi(a_i) - q_\xi(a_{i+1}) - 1$$

and

$$\tau_{t_i}(q_\xi)(b_{i+1}) = q_\xi(b_{i+1}) + (q_\xi(t_i) + 1) = q_\xi(b_{i+1}) - q_\xi(a_i) + q_\xi(a_{i+1}) + 1$$

which establishes the required formula. $\square$

Proof of Theorem 2.9. Consider the element

$$(A_1, B_1, \dots, A_g, B_g, R_1, \dots, R_b) \in \theta_r(\Sigma_{g,b+1}; \delta).$$

We will show how to reduce this to an element in standard form.

By iteratedly applying τ_{a_i} and τ_{b_i} we can reduce the pair (A_i, B_i) to the form $(-1, N)$. It is easy to check that the number $\gcd(A_i + 1, B_i + 1) \in \mathbb{Z}/r$ is invariant under τ_{a_i} and τ_{b_i} , so that $N = \gcd(A_i + 1, B_i + 1) - 1$; let us call this $G(A_i, B_i)$. It is not necessary for the proof, but to relate this to the function A we remark that $G(A_i, B_i) \equiv A_i B_i \pmod{2}$. Hence

$$(2.5) \quad (A_1, B_1, A_2, B_2, \dots, A_g, B_g, R_1, \dots, R_b)$$

is equivalent to

$$(2.6) \quad (-1, G(A_1, B_1), -1, G(A_2, B_2), \dots, -1, G(A_g, B_g), R_1, \dots, R_b).$$

Applying τ_{t_i} sends

$$(-1, G(A_i, B_i), -1, G(A_{i+1}, B_{i+1})) \text{ to } (-1, G(A_i, B_i) - 1, -1, G(A_{i+1}, B_{i+1}) + 1),$$

so applying it iteratedly to (2.6) gives

$$(2.7) \quad (-1, 0, -1, 0, \dots, -1, \sum_{i=1}^g G(A_i, B_i), R_1, \dots, R_b).$$

Let us write $N := \sum_{i=1}^g G(A_i, B_i)$ temporarily. We will now show how to reduce the numbers R_i to a more standard form. Define the simple closed curve u_i to be the connected sum of a_g and ∂_i , formed as indicated in Figure 1 to only intersect b_g , so $q_\xi(u_i) = A_g + \delta_i$. Twisting along u_1 gives

$$\tau_{u_1} \cdot (-1, 0, -1, 0, \dots, -1, N, R_1, \dots, R_r) = (-1, 0, -1, 0, \dots, -1, N + \delta_1, R_1 + \delta_1, \dots, R_r),$$

and twisting backwards around ∂_1 gives $(-1, 0, -1, 0, \dots, -1, N + \delta_1, R_1 - 1, \dots, R_r)$.

Repeating R_1 times gives $(-1, 0, -1, 0, \dots, -1, N + R_1 \delta_1, 0, R_2, \dots, R_r)$, and continuing in this way, by twisting around $u_2, u_3, \dots$ we can arrive at

$$(2.8) \quad (-1, 0, -1, 0, \dots, -1, \sum_{i=1}^g G(A_i, B_i) + \sum R_i \delta_i, 0, \dots, 0).$$

This much holds for $g \geq 1$.

If $g \geq 2$ we may apply the sequence

$$\begin{aligned} (-1, 0, -1, N) &\xrightarrow{\tau_{t_{g-1}}^2} (-1, -2, -1, N+2) \xrightarrow{\tau_{b_{g-1}}^2} (0, -2, -1, N+2) \\ &\xrightarrow{\tau_{a_{g-1}}^2} (0, 0, -1, N+2) \xrightarrow{\tau_{b_{g-1}}^2} (-1, 0, -1, N+2) \end{aligned}$$

to find that there are at *most* two orbits. If r is even the invariant A shows there are at *least* two orbits, and we are done. If r is odd note that we have put everything in the form $(-1, 0, \dots, -1, X, 0, \dots, 0)$ for $X \in \mathbb{Z}/r$, but this is equivalent to $(-1, 0, \dots, -1, X+2, 0, \dots, 0)$. Thus there is a single orbit in this case. $\square$

2.5. Gluing θ_r -surfaces. In this section we will discuss how the invariant A behaves with respect to gluing θ_r -surfaces. In order to do so effectively, it is convenient to discuss *connected cobordisms with θ_r -structure*. This simply means that we have designated incoming and outgoing boundaries, the marked points on the outgoing boundaries are framed using the boundary orientation and the inwards normal vector, and the marked points on the incoming boundaries are framed using the boundary orientation and the outwards normal vector. Furthermore, we have an ordering of first the outgoing boundaries and then the incoming boundaries, with respect to which we compute the generalised Arf invariant. We call this data a *structured θ_r -surface*.

Given cobordisms with boundary condition (Σ, δ) and (Σ', δ') , an identification $\psi : \partial_{out}\Sigma \cong \partial_{in}\Sigma'$ between incoming and outgoing and a choice of covering isomorphism $\phi : \delta|_{\partial_{out}\Sigma} \cong \delta'|_{\partial_{in}\Sigma'}$ between boundary conditions, determines a gluing map

$$(\psi, \phi) : \theta_r(\Sigma; \delta) \times \theta_r(\Sigma'; \delta') \longrightarrow \theta_r(\Sigma \cup_\psi \Sigma'; \delta \cup_\phi \delta')$$

which induces a map

$$(\psi, \phi) : \theta_r(\Sigma; \delta)/\Gamma^+(\Sigma) \times \theta_r(\Sigma'; \delta')/\Gamma^+(\Sigma) \longrightarrow \theta_r(\Sigma \cup_\psi \Sigma'; \delta \cup_\phi \delta')/\Gamma^+(\Sigma \cup_\psi \Sigma').$$

Under the identification of $\theta_r(\Sigma; \delta)/\Gamma^+(\Sigma)$ with $\pi_0(\mathcal{M}^{\theta_r}(\Sigma; \delta))$, this is nothing but the gluing map between these moduli spaces, at the level of π_0 .

We will determine the effect on this map on the invariant A . In order for this to be meaningful, we must declare how to impose the data of a structured θ_r -surface on $(\Sigma \cup_\psi \Sigma'; \delta \cup_\phi \delta')$: we use the marked points of the two surfaces and the induced ordering on the unglued boundaries.

Lemma 2.11. *Let r be even, and Σ and Σ' be connected. The function A evaluated on $(\psi, \phi)(\ell_\Sigma, \ell_{\Sigma'})$ is given by*

$$(2.9) \quad A(\ell_\Sigma) + A(\ell_{\Sigma'}) + \delta|_{\partial_0\Sigma}$$

Proof. This is immediate from the formula for A , by considering the new genus that may be formed in such a gluing. There are two important points:

- (i) When gluing two arcs together at a single end, the total monodromy along them adds, and so the value of the invariant adds, *but then one is added*.
- (ii) When two arcs are joined to create a new simple closed curve, the value of the invariant along it is the sum of the values along the two curves. In light of the previous point this is counterintuitive, but has to do with the canonical “straightening” done to arcs ending at the outgoing boundary or starting at the incoming boundary: the two half turns do not cancel, they *add*.

$\square$

In trying to understand this lemma, we remark that the reader should be aware that it is *only stated for connected surfaces*. We will discuss the failure of A to be either defined or to enjoy any good properties on non-connected surfaces elsewhere. The final tool we shall need is

Lemma 2.12. *In genus 1 the maps*

$$\theta_r(\Sigma_{1,b}; \delta) / \Gamma^+(\Sigma_{1,b}) \longrightarrow \theta_r(\Sigma_{1,b+1}; \delta') / \Gamma^+(\Sigma_{1,b+1})$$

that glue on a new boundary are surjective.

Proof. By the proof of Theorem 2.9, elements of $\theta_r(\Sigma_{1,b}; \delta') / \Gamma^+(\Sigma_{1,b})$ may be represented by $(-1, X, 0, \dots, 0)$. If we glue on the pair of pants represented by (R_1, R_2) along its last boundary to the first boundary of Σ , we obtain the element

$$(-1, X, R_2 + 1, \dots, R_2 + 1, R_1)$$

which may be reduced to $(-1, X + (R_2 + 1) \sum_{i=1}^{b-1} \delta_i + R_1 \delta'_1, 0, \dots, 0)$. By varying X we see that may obtain every element of the form $(-1, Y, 0, \dots, 0)$, as required. $\square$

2.6. Homological stability for framed and r -Spin surfaces. We will now describe how to use the above results to apply the main theorem of [18] to prove homological stability for framed and r -Spin surfaces. To do so we must necessarily use terminology introduced in [18], but we restrict its use to the proof. To be concise we give a single statement for all r : the statement for $r = 0$ is that for framed surfaces.

Theorem 2.13. *The moduli spaces of θ_r -surfaces exhibit homological stability. More precisely,*

- (i) *Any $\alpha(g) : \mathcal{M}^{\theta_r}(\Sigma_{g,b}) \rightarrow \mathcal{M}^{\theta_r}(\Sigma_{g+1,b-1})$ is a homology epimorphism in degrees $5* \leq 2g - 1$ and a homology isomorphism in degrees $5* \leq 2g - 6$.*
- (ii) *Any $\beta(g) : \mathcal{M}^{\theta_r}(\Sigma_{g,b}) \rightarrow \mathcal{M}^{\theta_r}(\Sigma_{g,b+1})$ is a homology epimorphism in degrees $5* \leq 2g - 2$ and a homology isomorphism in degrees $5* \leq 2g - 7$. If one of the created boundary conditions is trivial, it is a split homology monomorphism in all degrees.*
- (iii) *Any $\gamma(g) : \mathcal{M}^{\theta_r}(\Sigma_{g,b}) \rightarrow \mathcal{M}^{\theta_r}(\Sigma_{g,b-1})$ for $b \geq 2$ is a homology isomorphism in degrees $5* \leq 2g - 2$, and a split homology epimorphism in all degrees.*

Proof. In order to apply Theorem 7.1 of [18], we must verify two conditions: that θ_r -structures stabilise on π_0 at genus h , and that they are k -trivial. We claim that this is true for $(h, k) = (2, 4)$, then solving the recurrence relations of [18, §7.1] with this data will give the stated stability ranges.

The path components of the moduli spaces of θ_r -surfaces stabilise at genus 2 by Theorem 2.9 and Lemma 2.12. It follows from [18, Proposition 6.4] that θ_r -structures are 5-trivial, and from this we may already deduce homology stability in a certain range. However, we shall show that we in fact have 4-triviality, giving a better stability range.

To do this we refer the reader to the definition of 4-triviality in [18, §6.1]. For the case of α type stabilisation maps, we must show that given a diagram of θ_r -surfaces with compatible boundary conditions as shown in Figure 2, such that $\Sigma_L \circ \Sigma_T \cong \Sigma_B \circ \Sigma_R$ as θ_r -surfaces, there is a diagonal θ_r -surface Σ_Δ with boundary conditions $(D, A + B)$ such that $\Sigma_B \circ \Sigma_\Delta \cong \Sigma_L$ and $\Sigma_\Delta \circ \Sigma_T \cong \Sigma_R$.

By Lemma 2.12, the map

$$-\circ \Sigma_T : \theta_r(\Sigma_{1,1+2}; D + A \sqcup B) / \Gamma^+(\Sigma_{1,1+2}) \longrightarrow \theta_r(\Sigma_{1,2+2}; E \sqcup F + A \sqcup B) / \Gamma^+(\Sigma_{1,2+2})$$

is surjective, so let us choose Σ_Δ to be some preimage of Σ_R . By construction, $\Sigma_\Delta \circ \Sigma_T \cong \Sigma_R$, as required. We now wish to use the generalised Arf invariant, so must give the structure of a structured θ_r -surface on each cobordism: we choose




FIGURE 2. The given data to show 4-triviality with respect to α type stabilisation maps. The two θ_r -surfaces obtained by composition are isomorphic, and we must provide a genus 1 θ_r -surface from D to $A \sqcup B$ such that for each triangle the two obtained θ_r -surfaces are isomorphic.

a marked point on each boundary $A, B, \dots, F$, and choose an ordering of the boundary components of each cobordism, with all the outgoing boundaries first. In terms of Figure 2, let us order them clockwise starting at the rightmost outgoing boundary (so B is the first boundary component of Σ_R).

Applying the invariant A and the formula for gluing cobordisms to the given surfaces gives the equation

$$A(\Sigma_R) + A(\Sigma_B) + B = A(\Sigma_L) + A(\Sigma_T) + D \in \mathbb{Z}/2$$

and applying it to $\Sigma_\Delta \circ \Sigma_T \cong \Sigma_R$ gives the equation

$$A(\Sigma_R) = A(\Sigma_\Delta) + A(\Sigma_T) + D \in \mathbb{Z}/2.$$

Combining these gives that

$$A(\Sigma_B \circ \Sigma_\Delta) = A(\Sigma_\Delta) + A(\Sigma_B) + B = A(\Sigma_L).$$

As $\Sigma_B \circ \Sigma_\Delta$ and Σ_L are surfaces of genus 2, the invariant A is complete by Theorem 2.9 and hence $\Sigma_B \circ \Sigma_\Delta \cong \Sigma_L$ as θ_r -surfaces, as required. The argument for stabilisation maps of type β is the same. $\square$

2.7. Homological stability for closing the last boundary. Theorem 2.13 gives homology stability for almost all the stabilisation maps one might need, except for those which close off the last boundary. In the case of framings there are no such maps of interest: the only frameable closed surface has genus 1, for which the stability range is empty. Thus in this section we will restrict our attention to r -Spin structures only.

The paper [18] includes a general theorem on stability for closing the last boundary, whose hypotheses are that the relevant tangential structure has homological stability for all other stabilisation maps, and that it satisfies a condition called *closeability* [18, Definition 10.5].

Theorem 2.14. *Any $\gamma(g) : \mathcal{M}^{\text{Spin}^r}(\Sigma_{g,1}; \delta) \rightarrow \mathcal{M}^{\text{Spin}^r}(\Sigma_g)$ is a homology isomorphism in degrees $5* \leq 2g - 2$, and a homology epimorphism in degrees $5* \leq 2g + 3$.*

Proof. We will verify the conditions for closeability for the tangential structures θ_r , $r \geq 1$, and so assume the language of [18]. The natural map $B_0^{\theta_r} \rightarrow B_0^{SO}$, from the space of r -Spin structures on D^2 to the space of orientations of D^2 , induces a bijection on π_0 as the fibre of $\theta_r : B\text{Spin}^r(2) \rightarrow BSO(2)$ is path-connected.

Hence $\pi_0(C_0^{\theta_r}(\Sigma)) \cong \pi_0(C_0^{SO}(\Sigma))$ for any orientable surface Σ , and we showed in [18, §10.3.1] that orientations are a closeable tangential structure, so θ_r -structures are also. The theorem now follows from [18, Theorem 10.6]. $\square$

3. APPLICATIONS OF HOMOLOGY STABILITY FOR FRAMED SURFACES

The methods of Galatius, Madsen, Tillmann and Weiss [8, Section 7] take the homological stability theorem for framed surfaces and imply a homology equivalence

$$\text{colim}_{g \rightarrow \infty} \mathcal{M}^{\text{fr}}(\Sigma_{g,1+1}) =: \mathcal{M}^{\text{fr}}(\Sigma_{\infty}) \longrightarrow \Omega^{\infty} \mathbf{S}^{-2} \simeq \Omega^2 Q(S^0)$$

where the colimit is formed by gluing on framed tori with two boundary components.

Corollary 3.1. *The moduli space of framed surfaces $\mathcal{M}^{\text{fr}}(\Sigma_{g,b}; \delta)$ has the integral homology of $\Omega^2 Q(S^0)$ in degrees $5* \leq 2g - 7$. In particular it has trivial rational homology in these degrees. The same is true for the framed mapping class group $\Gamma^{\text{fr}}(\Sigma_{g,b}; \xi)$.*

Corollary 3.2. *The abelianisation of $\Gamma^{\text{fr}}(\Sigma_{g,b}; \xi)$ is $\mathbb{Z}/24$, as long as $g \geq 6$.*

Proof. The abelianisation is simply the first integral homology of this group, which for $g \geq 6$ coincides with the first integral homology of $\Omega^{\infty} \mathbf{S}^{-2}$. By Hurewicz' theorem this is the same as $\pi_1(\Omega^{\infty} \mathbf{S}^{-2}) = \pi_3^s$, the third stable stem, which is well known to be cyclic of order 24. $\square$

A diffeomorphism $\varphi : \Sigma_{g,1} \rightarrow \Sigma_{g,1}$ that preserves a framing ξ (up to isomorphism) represents an element $[\varphi] \in \Gamma^{\text{fr}}(\Sigma_{g,1}; \xi)$ in the framed mapping class group, and one may ask what class it represents in the abelianisation $\mathbb{Z}/24$.

The mapping torus $\Sigma_{g,1} \rightarrow M_{\varphi} \rightarrow S^1$ is a smooth 3-manifold with boundary $\partial \Sigma_{g,1} \times S^1$, and we may frame M_{φ} by choosing to take the Lie framing of S^1 and the framing ξ along the fibres. On the boundary this framing of $\partial \Sigma_g^1 \times S^1$ bounds a framed solid torus by filling in a trivially framed disc in the S^1 direction. Gluing such a framed solid torus in we obtain a closed framed 3-manifold M^3 , which represents an element of $\pi_3^s \cong \mathbb{Z}/24$.

It is well known that the element of $\mathbb{Z}/24$ represented by a framed 3-manifold M may be computed as follows. M is in particular a Spin 3-manifold, and any closed Spin 3-manifold bounds a Spin 4-manifold, so let $M^3 = \partial W^4$. The first Pontrjagin class of W may be represented by a cocycle which is identically zero on $\partial W = M$ as this is framed, so gives an element $p_1 \in H^4(W, M; \mathbb{Z})$. We may then form

$$-\frac{1}{48} \langle p_1, [W, M] \rangle \in \mathbb{Q}.$$

A different choice W' of Spin manifold means that $W \cup W'$ is a closed Spin manifold, and the two invariants defined above differ by $-\frac{1}{48} \langle p_1, [W \cup W'] \rangle$, which is an integer (as it is half the $\hat{A}$ -genus of the closed Spin manifold $W \cup W'$, which is even by Rokhlin's theorem). Thus we obtain a well defined element of $\mathbb{Q}/\mathbb{Z}$ which lies in the subgroup generated by $2/48$, which is isomorphic to $\mathbb{Z}/24$. Hence, in principle we have described the map $\Gamma^{\text{fr}}(\Sigma_{g,1}; \xi)$, though in practice it is a matter of some difficulty to evaluate the above characteristic number.

4. THE MODULI SPACES OF $\text{Pin}^\pm$ SURFACES

In the introduction we described how to a map $\theta : \mathbf{X} \rightarrow BO(2)$ we can associate a moduli space of surfaces with θ -structure, so to define the moduli spaces of Pin -surfaces it is enough to describe this map.

This is complicated by the fact that there are two forms of $\text{Pin}(2)$ group, $\text{Pin}^+(2)$ and $\text{Pin}^-(2)$, which are most easily distinguished by saying that they are the central extensions of $O(2)$ classified by the classes w_2 and $w_2 + w_1^2$ in $H^2(BO(2); \mathbb{Z}/2)$ respectively (which we identify with the continuous group cohomology of $O(2)$). The classifying spaces of both groups fit into principal homotopy fibre sequences

$$(4.1) \quad B\mathbb{Z}/2 \longrightarrow B\text{Pin}^\pm(2) \xrightarrow{\theta^\pm} BO(2),$$

and we define $\theta^\pm$ to be the indicated map in the defining extension. Let $\gamma_2^{\text{Pin}^\pm} \rightarrow B\text{Pin}^\pm(2)$ and $\gamma_2 \rightarrow BO(2)$ denote the tautological bundles.

Definition 4.1. For each choice of sign, the *moduli space of $\text{Pin}^\pm$ surfaces* of topological type F and boundary condition $\delta : TF|_{\partial F} \rightarrow \gamma_2^{\text{Pin}^\pm}$, denoted $\mathcal{M}^{\text{Pin}^\pm}(F; \delta)$, is that associated to the map $\theta^\pm$.

There is a natural fibration

$$\text{Bun}_\partial(TF, \gamma_2^{\text{Pin}^\pm}; \delta) \longrightarrow \text{Bun}_\partial(TF, \gamma_2; [\delta]),$$

where $[\delta]$ denotes the composition $TF|_{\partial F} \xrightarrow{\delta} \gamma_2^{\text{Pin}^\pm} \rightarrow \gamma_2$. The fibres of this map are either empty (if F does not admit a $\text{Pin}^\pm$ -structure compatible with δ), or else are homotopy equivalent to $\text{map}_*(F/\partial F, B\mathbb{Z}/2)$, where we consider $F/\partial F$ to be F with a disjoint basepoint added if $\partial F = \emptyset$.

Hence if $\mathcal{M}^{\text{Pin}^\pm}(F; \delta)$ is non-empty there is a fibration sequence

$$(4.2) \quad \text{map}_*(F/\partial F, B\mathbb{Z}/2) \longrightarrow \mathcal{M}^{\text{Pin}^\pm}(F; \delta) \longrightarrow \mathcal{M}^{O(2)}(F; [\delta]),$$

obtained by taking the homotopy quotient by $\text{Diff}_\partial(F)$.

Remark 4.2. Both $\text{Pin}^+(2)$ and $\text{Pin}^-(2)$ become isomorphic to $\text{Spin}(2)$ when restricted to covering groups of $SO(2)$. Hence if a surface is orientable and has boundary, a $\text{Pin}^\pm$ -structure on it is nothing but a (2-)Spin structure. As we have dealt with this situation in the previous sections, from now on we will assume that we only consider non-orientable surfaces.

4.1. $\text{Pin}^\pm$ mapping class groups. We consider the long exact sequence on homotopy groups associated to the fibration (4.2). If ∂F is non-empty, the fibre is homotopy-discrete so this induces an exact sequence on homotopy groups


$$0 \longrightarrow \Gamma^{\text{Pin}^\pm}(F; \xi) \longrightarrow \Gamma(F) \longrightarrow H^1(F, \partial F; \mathbb{Z}/2) \longrightarrow \pi_0(\mathcal{M}^{\text{Pin}^\pm}(F; \delta)) \longrightarrow *$$

where $\Gamma^{\text{Pin}^\pm}(F; \xi) := \pi_1(\mathcal{M}^{\text{Pin}^\pm}(F; \delta), \xi)$. This identifies the $\text{Pin}^\pm$ mapping class group $\Gamma^{\text{Pin}^\pm}(F; \xi)$ as the subgroup of the unoriented mapping class group consisting of those diffeomorphisms which fix the $\text{Pin}^\pm$ -structure ξ up to isomorphism.

If ∂F is empty, there is an exact sequence

$$0 \longrightarrow \mathbb{Z}/2 \longrightarrow \Gamma^{\text{Pin}^\pm}(F; \xi) \longrightarrow \Gamma(F) \longrightarrow \dots$$

which identifies the $\text{Pin}^\pm$ mapping class group with an extension by $\mathbb{Z}/2$ of the subgroup of the unoriented mapping class group of elements fixing the $\text{Pin}^\pm$ -structure ξ up to isomorphism. This is analogous to the 2-Spin mapping class group in the case of oriented surfaces.

FIGURE 3. Curves and arcs on $S_{n,b+1}$.

4.2. The set of $\text{Pin}^\pm$ -structures. In order to apply the results of [18], we must compute the sets of components $\pi_0(\mathcal{M}^{\text{Pin}^\pm}(F; \delta))$, at least when the genus of F is large. This is the same as the quotient of the set of $\text{Pin}^\pm$ -structures

$$\text{Pin}^\pm(F, \delta) := \pi_0(\text{Bun}_\partial(TF, \gamma_2^{\text{Pin}^\pm}; \delta)),$$

by the action of the unoriented mapping class group $\Gamma(F)$.

Remark 4.3. By considering the fibration

$$\text{map}_*(F/\partial F, B\mathbb{Z}/2) \longrightarrow \text{Bun}_\partial(TF, \gamma_2^{\text{Pin}^\pm}; \delta) \longrightarrow \text{Bun}_\partial(TF, \gamma_2; [\delta])$$

when $\text{Bun}_\partial(TF, \gamma_2^{\text{Pin}^\pm}; \delta)$ is non-empty, we see that the set $\text{Pin}^\pm(F, \delta)$ is a torsor for $H^1(F, \partial F; \mathbb{Z}/2)$.

As in the r -Spin case, we choose once and for a $\text{Pin}^\pm$ -structure on $\mathbb{R}^2$, which induces a *standard* $\text{Pin}^\pm$ -structure on any framed rank 2 vector bundle. On the non-orientable surface $S_{n,b+1}$ having some $\text{Pin}^\pm$ -structure ξ with boundary condition δ , we choose a marked point on each boundary component, which has a canonical framing coming from the orientation of the boundary and the inwards pointing normal vector. After perhaps changing δ to an isomorphic boundary condition, we may assume that it is standard at each marked point with respect to this framing.

Let $\{a_1, \dots, a_n, \partial_0, \dots, \partial_b\}$ be the collection of simple closed curves as shown in Figure 3, and $\{r_1, \dots, r_b\}$ be the collection of simple arcs. We also write $a_i + a_j$ and $a_i + a_j + a_k + a_l$ for the simple closed curves shown, indicating the homology class they represent. There is a presentation of the first integral homology of $S_{n,b+1}$ as

$$H_1(S_{n,b+1}; \mathbb{Z}) = \mathbb{Z}\langle a_1, \dots, a_n, \partial_0, \dots, \partial_b \rangle / \langle \partial_0 + \dots + \partial_b + 2(a_1 + \dots + a_n) \rangle.$$

Definition 4.4. Given a $\text{Pin}^\pm$ -structure ξ on F , along any simple closed curve a or simple arc r between standard marked points we have a canonical reduction of structure group from $\text{Pin}^\pm(2)$ to $\text{Pin}^\pm(1) = O(1) \times \mathbb{Z}/2$. The first factor detects orientability of the loop or arc. Taking the value of the monodromy along a or r on the second factor and adding 1 defines $q_\xi(a)$ or $q_\xi(r) \in \mathbb{Z}/2$.

Given a Pin^- -structure ξ on F , along any simple closed curve a or simple arc r between standard marked points we have a canonical reduction of structure group

from $\text{Pin}^-(2)$ to $\text{Pin}^-(1) = \mathbb{Z}/4$. Adding 2 to the monodromy along a or r defines $q_\xi(a)$ or $q_\xi(r) \in \mathbb{Z}/4$. The reduction of $q_\xi(x)$ modulo 2 detects orientability of the loop or arc.

We write $\text{Mon}(a)$ or $\text{Mon}(r)$ for the monodromy (in $\mathbb{Z}/2$ or $\mathbb{Z}/4$) around a curve a or arc r between standard marked points.

The convention of adding the monodromy around a disc to the actual monodromy in this definition is as a normalisation: now $q_\xi(a) = 0$ if and only if a bounds a disc with $\text{Pin}^\pm$ -structure. We leave the proof of the fact that the monodromy around a disc is 1 in the Pin^+ case and 2 in the Pin^- case to the reader: it is an interesting exercise in characteristic classes.

Remark 4.5 (Boundary conditions). For each oriented boundary component ∂_i of a surface F with $\text{Pin}^\pm$ -structure ξ , we may evaluate q_ξ on the simple closed curve ∂_i to get δ_i . For Pin^+ -structures this lies in $\mathbb{Z}/2$ and for Pin^- -structures it lies in $2\mathbb{Z}/4 = \mathbb{Z}/2$ as the curve is orientation-preserving. This allows us to identify boundary conditions on $S_{n,b+1}$ for either of these tangential structures with elements of $(\mathbb{Z}/2)^{b+1}$, and $\delta_i = 0$ precisely when this boundary component bounds a disc.

There is a map $e^+ : \text{Pin}^+(S_{n,b+1}; \delta) \rightarrow (\mathbb{Z}/2)^{n+b}$ (or to $(\mathbb{Z}/2)^n$ if the surface is closed) given by $e^+(\xi) = (q_\xi(a_1), \dots, q_\xi(a_n), q_\xi(r_1), \dots, q_\xi(r_b))$. Similarly, there is a map by evaluation on the a_i and r_i .

Proposition 4.6. *If the set $\text{Pin}^+(S_{n,b+1}; \delta)$ is non-empty, e^+ is a bijection. If the set $\text{Pin}^-(S_{n,b+1}; \delta)$ is non-empty, e^- is a bijection onto the subset $\{[1], [3]\}^n \times \{[0], [2]\}^b$.*

Proof. Certainly both maps are injective: once a $\text{Pin}^\pm$ -structure ξ is determined on $a_1, \dots, a_n, r_1, \dots, r_b$ and the boundary by δ , it remains to give a $\text{Pin}^\pm$ -structure on a disc satisfying a certain boundary condition (which depends only on δ). If this is possible, it is possible in at most one way, as the space of $\text{Pin}^\pm$ -structures on the disc is either empty or a torsor for $\Omega^2 B\mathbb{Z}/2 \simeq *$.

The image of e^- certainly lies in the subset $\{[1], [3]\}^n \times \{[0], [2]\}^b$ due to the values a Pin^- -structure may take around orientation-reversing and -preserving arcs. Now note that $\text{Pin}^\pm(F; \delta)$ is either empty or a $H^1(F, \partial F; \mathbb{Z}/2)$ -torsor, so counting now implies that the maps are bijections. $\square$

Remark 4.7. We should mention to what extent the functions e^+ and e^- are unique. We have only defined them for boundary conditions with a choice of standard marked point on each boundary, and an ordering of the boundaries. Varying either of these choices will vary the functions.

Proposition 4.8. *Let $\xi \in \text{Pin}^\pm(F; \delta)$ be a $\text{Pin}^\pm$ -structure and $g \in H^1(F, \partial F; \mathbb{Z}/2)$ be a cohomology class. The torsor structure gives a new $\text{Pin}^\pm$ -structure $g \cdot \xi$ and on a simple closed curve or simple closed arc a we have the formula*

$$q_{g \cdot \xi}(a) = q_\xi(a) + g(a) \cdot \text{Mon}(D^2),$$

where $\text{Mon}(D^2)$ is 1 in the Pin^+ case and 2 in the Pin^- case.

Proof. Direct from the definition of the torsor structure. $\square$

Using Proposition 4.6, we may study the action of $\Gamma(S_{n,b+1})$ on $\text{Pin}^\pm(F; \delta)$ by studying its action in the ‘‘coordinate system’’ given by $e^\pm$. To do so we will require various formulæ for computing with q_ξ .

Lemma 4.9. *Let $S_{n,1}$ be a non-orientable surface with $\text{Pin}^\pm$ -structure ξ . Then $\delta_0 = 0$ for Pin^- , and $\delta_0 = n \bmod 2$ for Pin^+ . Let $\Sigma_{g,1}$ be an orientable surface with $\text{Pin}^\pm$ -structure ξ . Then $\delta_0 = 0$ for both Pin^- and Pin^+ .*

More generally, for a surface with multiple boundaries the same is true for $\sum \delta_i$.

Proof. The value of δ_0 is the value of $w_2 + w_1^2$ or w_2 respectively evaluated on S_n or Σ_g . This is 0 and $n \bmod 2$ respectively on S_n , and always 0 on Σ_g . $\square$

Lemma 4.10. *If a and b are disjoint simple closed curves (or a simple closed and a simple arc), and $a\#b$ is their oriented connected sum, then*

$$q_\xi(a\#b) = q_\xi(a) + q_\xi(b).$$

Proof. We may suppose that the connect sum is formed in a small disc on the surface where we have trivialised the $\text{Pin}^\pm$ -structure. Hence the difference in total monodromies of $a\sqcup b$ and $a\#b$ is the monodromy around a small disc, and the claim follows as q_ξ is corrected from the actual monodromy by precisely the monodromy around a small disc. $\square$

Lemma 4.11. *Let a and b be simple closed curves (or a simple closed and a simple arc), and c be a simple closed curve (or a simple closed arc).*

(i) *For Pin^- , if the $\mathbb{Z}/2$ -homology classes satisfy $[c] = [a] + [b]$, then*

$$q_\xi(c) = q_\xi(a) + q_\xi(b) + 2 \cdot \langle a, b \rangle \in \mathbb{Z}/4.$$

(ii) *For Pin^+ , if the $\mathbb{Z}/4$ -homology classes satisfy $[c] = [a] + [b]$, then*

$$q_\xi(c) = q_\xi(a) + q_\xi(b) + \langle a, b \rangle \in \mathbb{Z}/2.$$

If b bounds a disc, we are asserting that $q_\xi(a)$ only depends on the ($\mathbb{Z}/2$ or $\mathbb{Z}/4$, respectively) homology class of a .

Proof. Firstly, we perturb a and b so that they cross transversely. Near each intersection point we can cut out the intersection and glue the four incoming arcs together in pairs, to get a homologous collection of N disjoint simple closed curves (with perhaps a single arc). Each time we do this, the total monodromy around all these curves does not change, so we still have total monodromy $Mon(a) + Mon(b)$. When we connect-sum together the N disjoint curves, we end up with a single curve with monodromy

$$Mon(a) + Mon(b) + Mon(D^2) \cdot (N - 1),$$

as in the proof of the lemma above, and hence a single curve with invariant

$$q_\xi(a) + q_\xi(b) + Mon(D^2) \cdot N.$$

We now simply remark that during the process of eliminating intersection points, the value of

$$\#\{\text{components}\} + \#\{\text{intersection points}\} \in \mathbb{Z}/2$$

is constant, so in particular $N \equiv \langle a, b \rangle \bmod 2$, and the curve we have constructed has invariant $q_\xi(a) + q_\xi(b) + Mon(D^2) \cdot \langle a, b \rangle$.

It is now enough to prove the homology invariance of q_ξ : if a and c are ($\mathbb{Z}/4$ or $\mathbb{Z}/2$, respectively) homologous, then $q_\xi(a) = q_\xi(c)$. By taking the difference of a and c , and performing the above manouver to get a single simple closed curve, it is enough to show that if a is homologically trivial then $q_\xi(a) = 0$.

To prove this we use the classification of simple closed curves in a surface (which is of course a simple consequence of the classification of surfaces). For the Pin^- case, a simple closed curve on a non-orientable surface can be trivial in $\mathbb{Z}/2$ -homology only if it bounds a subsurface, then by Lemma 4.9 the value of the invariant on it is zero. For the Pin^+ case, a simple closed curve on a non-orientable surface can be

trivial in $\mathbb{Z}/4$ -homology only if it bounds an orientable subsurface, then by Lemma 4.9 the value of the invariant is zero. $\square$

We learnt the above results from the work of Degtyarev–Finashin [4], but have adapted the results and proofs in the form that is most convenient to us. Using the above lemma one can extend q_ξ uniquely to a function on the first $\mathbb{Z}/2$ - or $\mathbb{Z}/4$ -homology of $S_{n,b+1}$, and the lemma shows that q_ξ is a quadratic refinement of the intersection form.

We wish to compute the action of the unoriented mapping class group $\Gamma(S_{n,b+1})$ on the sets $\text{Pin}^\pm(S_{n,b+1}; \delta)$. Recall that the unoriented mapping class group is generated by Dehn twists and crosscap slides [13]. By the following lemma, it will be enough to consider just Dehn twists.

Lemma 4.12. *Crosscap slides act trivially on $\text{Pin}^\pm(S_{n,b+1}; \delta)$.*

Proof. Crosscap slides act trivially on the integral homology of $S_{n,b+1}$. If τ is such a diffeomorphism then

$$q_{\tau^*\xi}(a) = \tau^*q_\xi(a) = q_\xi(\tau(a)) = q_\xi(a)$$

as $\tau(a)$ is homologous to a . $\square$

Hence we only need to study the action of Dehn twists on the set of $\text{Pin}^\pm$ -structures. We will require the following formulæ, where a is a simple closed curve, τ_a is the Dehn twist around a , and x is a simple arc or simple closed curve: for Pin^+ -structures

$$q_{\tau_a^*\xi}(x) = q_\xi(x) + \langle a, x \rangle (q_\xi(a) - 1) \in \mathbb{Z}/2,$$

and for Pin^- -structures

$$q_{\tau_a^*\xi}(x) = q_\xi(x) + \langle a, x \rangle (q_\xi(a) - 2) \in \mathbb{Z}/4.$$

Both equations come from the standard action of Dehn twists on homology, $\tau_a(x) = x + \langle a, x \rangle \cdot a$, and Lemma 4.11.

4.3. Diffeomorphism classes of Pin^+ -structures. Once we have a bijection $e^+ : \text{Pin}^+(S_{n,b+1}; \delta) \rightarrow (\mathbb{Z}/2)^{n+b}$, we define a function

$$A : \text{Pin}^+(S_{n,b+1}; \delta) \longrightarrow \mathbb{Z}/2$$

by the formula $\xi \mapsto \sum_{i=1}^n q_\xi(a_i)$.

Proposition 4.13. *The function A is $\Gamma(S_{n,b+1})$ -invariant.*

Proof. It is enough to show that it is invariant under Dehn twists. Let a be an orientation preserving simple closed curve represented in homology by $\sum A_i a_i + \sum \lambda_j \partial_j \in H_1(S_{n,b+1}; \mathbb{Z})$, so $\langle a, a \rangle = \sum A_i = 0$. We may then compute

$$\sum_{i=1}^n q_{\tau_a^*\xi}(a_i) = \sum_{i=1}^n q_\xi(a_i + \langle a_i, a \rangle \cdot a) = \sum_{i=1}^n q_\xi(a_i) + \sum_{i=1}^n A_i (1 + q_\xi(a)) = \sum_{i=1}^n q_\xi(a_i) \in \mathbb{Z}/2$$

so A is a diffeomorphism invariant of the Pin^+ -structure ξ . $\square$

Proposition 4.14. *The induced map $A : \text{Pin}^+(S_{n,b+1}; \delta) / \Gamma(S_{n,b+1}) \rightarrow \mathbb{Z}/2$ is a bijection if the source is non-empty and $n \geq 3$. It is a surjection for $n \geq 1$.*

Proof. We use the bijection e^+ to identify $\text{Pin}^+(S_{n,b+1}; \delta)$ with $\mathbb{Z}/2^{n+b}$, and write elements as $(A_1, \dots, A_n, R_1, \dots, R_b)$. If we take the simple closed curve $a_i + a_j$ from Figure 3 (when $n \geq 2$), we have

$$q_{\tau_{a_i+a_j}^*\xi}(a_s) = q_\xi(a_s + \langle a_i + a_j, a_s \rangle \cdot (a_i + a_j)) = \begin{cases} q_\xi(a_s) & s \neq i, j \\ 1 + q_\xi(a_j) & s = i \\ 1 + q_\xi(a_i) & s = j. \end{cases}$$

Thus given an element $(A_1, \dots, A_n) \in (\mathbb{Z}/2)^n$, we may permute a pair of entries and add 1 to both, via Dehn twists. In particular, we may remove pairs of 1's, and we may move a single 1 to be at any position (if $n \geq 3$). Thus we may reduce any element to the form

$$(0, 1, 0, \dots, 0, R_1, \dots, R_b) \text{ or } (0, 1, 1, 0, \dots, 0, R_1, \dots, R_b)$$

as long as $n \geq 3$.

Suppose that $R_i = 1$. If $\delta_i = 0$ then twisting along ∂_i gives

$$q_{\tau_{\partial_i}^*} \xi(r_i) = q_\xi(r_i) + q_\xi(\partial_i) - 1 = 0$$

and does not affect the other coordinates. If $\delta_i = 1$ then the simple closed curve $a_1 + a_2$ may be connect-summed to the simple closed curve ∂_i to give a curve x which intersects a_1 , a_2 and r_i once, the other curves and arcs not at all, and has $q_\xi(x) = 0$. Then

$$q_{\tau_x^*} \xi(r_i) = q_\xi(r_i) + q_\xi(x) - 1 = 0$$

so twisting around this sets R_i to 0. Doing this for each i allows us to reduce to an element of the form

$$(0, 1, 0, \dots, 0) \text{ or } (0, 1, 1, 0, \dots, 0)$$

as long as $n \geq 3$, so there are at most two orbits. On the other hand, the invariant A distinguishes these two elements, so there are precisely two orbits, distinguished by A . $\square$

We also make the following observation about the behaviour of A with respect to the $H^1(F, \partial F; \mathbb{Z}/2)$ -torsor structure on $\text{Pin}^+(F; \delta)$, which is immediate from the formula for A ,

$$(4.3) \quad A(g \cdot \xi) = A(\xi) + g \left(\sum_{i=1}^n a_i \right).$$

4.4. Diffeomorphism classes of Pin^- -structures. Once we have the map $e^- : \text{Pin}^-(S_{n,b+1}; \delta) \rightarrow (\mathbb{Z}/4)^{n+b}$ which is a bijection onto the subset $\{[1], [3]\}^n \times \{[0], [2]\}^b$, we define a function

$$A : \text{Pin}^-(S_{n,b+1}; \delta) \longrightarrow \mathbb{Z}/4$$

by the formula

$$\xi \mapsto \#\{q_\xi(a_i) = 1\} - \sum_{j=1}^b \delta_j \cdot \frac{q_\xi(r_j)}{2},$$

bearing in mind that r_i is an orientation-preserving arc so $q_\xi(r_i)$ is 0 or 2, and $\frac{q_\xi(r_i)}{2}$ is then defined to be 0 or 1.

Proposition 4.15. *The function A is $\Gamma(S_{n,b+1})$ -invariant.*

Proof. We only need to show that it is invariant under Dehn twists. Let x be an orientation-preserving simple closed curve represented in homology by $\sum X_i a_i + \sum \lambda_j \partial_j$, and ξ a Pin^- -structure. Then

$$q_{\tau_x^*} \xi(a_i) = q_\xi(a_i) + X_i(q_\xi(x) - 2)$$

and

$$q_{\tau_x^*} \xi(r_j) = q_\xi(r_j) + (\lambda_j - \lambda_0)(q_\xi(x) - 2)$$

so if $q_\xi(x) = 2$ the proposed invariant is trivially preserved. If $q_\xi(x) = 0$ then

$$\sum \delta_i q_{\tau_x^*} \xi(r_i)/2 - \sum \delta_i q_\xi(r_i)/2 = - \sum \delta_j (\lambda_j - \lambda_0).$$

Using Lemma 4.9, $\sum \delta_i = 0$ and so

$$-\sum_{j=1}^r \delta_j (\lambda_j - \lambda_0) = -\sum_{j=0}^r \delta_j \lambda_j = -q_\xi(x) + \sum X_i q_{\tau_x^* \xi}(a_i) = \sum X_i q_{\tau_x^* \xi}(a_i),$$

which is $\sum X_i (q_\xi(a_i) - 2X_i) = \sum X_i q_\xi(a_i) - 2\langle x, x \rangle = \sum X_i q_\xi(a_i)$, as $\langle x, x \rangle = 0$ because x is orientation-preserving.

On the other hand, $q_{\tau_x^* \xi}(a_i) = 1$ if and only if $q_\xi(a_i) = 1 + 2X_i$, which falls into two disjoint cases:

- (i) either $q_\xi(a_i) = 1$ and $X_i = 0$,
- (ii) or $q_\xi(a_i) = 3$ and $X_i = 1$.

Thus the difference $\#\{q_\xi(a_i) = 1\} - \#\{q_\xi(a_i) = 1 + 2X_i\}$ is the same as the difference $\#\{q_\xi(a_i) = 1, X_i = 1\} - \#\{q_\xi(a_i) = 3, X_i = 1\}$ which is $\sum X_i q_\xi(a_i)$, as required. $\square$

Proposition 4.16. *The induced map $A : \text{Pin}^-(S_{n,b+1}; \delta) / \Gamma(S_{n,b+1}) \rightarrow \mathbb{Z}/4$ is a bijection if the source is non-empty and $n \geq 3$.*

Proof. Let us use the bijection e^- to identify $\text{Pin}^-(S_{n,b+1}; \delta)$ with $\{[1], [3]\}^n \times \{[0], [2]\}^b$, and write elements as $(A_1, \dots, A_n, R_1, \dots, R_b)$. Note that

$$q_{\tau_{a_i+a_j}^* \xi}(a_s) = q_\xi(a_s + \langle a_i + a_j, a_s \rangle \cdot (a_i + a_j)) = \begin{cases} q_\xi(a_s) & s \neq i, j \\ q_\xi(a_j) & s = i \\ q_\xi(a_i) & s = j. \end{cases}$$

Thus given an element $(A_1, \dots, A_n) \in (1 + 2\mathbb{Z}/4)^n$ there is a diffeomorphism which permutes its entries arbitrarily. Note also that

$$q_{\tau_{a_i+a_j+a_k+a_l}^* \xi}(a_s) = \begin{cases} q_\xi(a_s) & i \neq i, j, k, l \\ q_\xi(a_j + a_k + a_l) & s = i \\ \text{etc.} & \end{cases}$$

This allows us to replace an occurrence of $(3, 3, 3, 3)$ in $(A_1, \dots, A_n)$ by $(1, 1, 1, 1)$.

Suppose that $R_i = 2$. If $\delta_i = 0$ then twisting along ∂_i sets R_i to 0. As $n \geq 3$ there is a pair of basis elements a_j, a_k such that $\{A_j, A_k\} = \{1, 1\}$ or $\{3, 3\}$. If $\delta_i = 2$ the simple closed curve $a_j + a_k$ may be connect-summed to the simple closed curve ∂_i to give a simple closed curve x which intersects a_j, a_k and r_i once, the other curves not at all, and has $q_\xi(x) = q_\xi(a_j) + q_\xi(a_k) + \delta_i = 0$. Then

$$q_{\tau_x^* \xi}(r_i) = q_\xi(r_i) + q_\xi(x) - 2 = 0$$

so twisting around this sets R_i to 0. This shows how to reduce to $R_1 = \dots = R_r = 0$. Thus for $n \geq 3$ there are at most 4 orbits. On the other hand, the invariant A distinguishes the four elements

$$(3, \dots, 3, 0, \dots, 0), (1, \dots, 3, \dots, 3, 0, \dots, 0), \dots, (1, 1, 1, 3, \dots, 3, 0, \dots, 0)$$

and so there are precisely four orbits, distinguished by A . $\square$

We also make the following observation about the behaviour of A with respect to the $H^1(F, \partial F; \mathbb{Z}/2)$ -torsor structure on $\text{Pin}^-(F; \delta)$, which is immediate from the formula for A ,

$$(4.4) \quad A(g \cdot \xi) = A(\xi) + \sum_{i=1}^n g(a_i) \cdot (q_\xi(a_i) - 2) - \sum_{j=1}^b \delta_j g(r_j).$$

4.5. Gluing $\text{Pin}^\pm$ surfaces. As in §2.5, we require a formula for the effect on the invariant A of gluing together surfaces. For our purposes we only need such a formula for gluing together cobordisms of the form $S_{n,1+1}$, and so we shall not investigate the existence of a formula for gluing more general cobordisms.

If (F, δ) and (F', δ') are cobordisms with $\text{Pin}^\pm$ boundary conditions and there is an identification $\psi : \partial_{in} F' \cong \partial_{out} F$ of boundary components along with a covering isomorphism $\phi : \delta|_{\partial_{in} F'} \cong \delta'|_{\partial_{out} F}$ between boundary conditions, we obtain a gluing map

$$\begin{aligned} (\psi, \phi) : \text{Pin}^\pm(F; \delta)/\Gamma(F) \times \text{Pin}^\pm(F'; \delta')/\Gamma(F') \\ \longrightarrow \text{Pin}^\pm(F \cup_\psi F'; \delta_{in} + \delta'_{out})/\Gamma(F \cup_\psi F'). \end{aligned}$$

We always suppose that each boundary components of each surface has a marked point with standard $\text{Pin}^\pm$ -structure, and that these are identified under (ψ, ϕ) .

Lemma 4.17. *Consider gluing together two $\text{Pin}^\pm$ cobordisms $S_{n,1+1}$ and $S_{n',1+1}$. In the Pin^+ case, the function A evaluated on the glued surface $(\psi, \phi)(\xi, \xi')$ is $A(\xi) + A(\xi')$. In the Pin^- case, the function A evaluated on the glued surface $(\psi, \phi)(\xi, \xi')$ is $A(\xi) + A(\xi') + \delta|_{\partial_{in} S_{n',1+1}}$.*

Proof. These formulas are immediate from the definitions of the invariant A , and in the Pin^- case the observation that when gluing two arcs together at a single end the monodromy adds, so the value of q_ξ adds, but then two is added. $\square$

We also require the following lemma concerning more general gluings, though not any explicit formula for the effect on the invariant A .

Lemma 4.18. *If we fix a (possibly orientable) $\text{Pin}^\pm$ -surface F' , then the map*

$$(\psi, \phi) : \text{Pin}^\pm(F; \delta)/\Gamma(F) \longrightarrow \text{Pin}^\pm(F \cup_\psi F'; \delta_{in} + \delta'_{out})/\Gamma(F \cup_\psi F')$$

is a bijection as long as F is non-orientable of genus ≥ 3 .

Proof. Note both sides have the same cardinality (2 in the Pin^+ case and 4 in the Pin^- case), and so it is enough to show that the map is surjective. For this we consider the commutative square

$$\begin{array}{ccc} \text{Pin}^\pm(F) & \xrightarrow{A} & \mathbb{Z}/2 \text{ or } \mathbb{Z}/4 \\ (\psi, \phi) \downarrow & & \downarrow \text{induced} \\ \text{Pin}^\pm(F \cup_\psi F') & \xrightarrow{A} & \mathbb{Z}/2 \text{ or } \mathbb{Z}/4 \end{array}$$

where the left vertical map is a map of torsors over the map

$$H^1(F, \partial F; \mathbb{Z}/2) \longrightarrow H^1(F \cup_\psi F', \partial(F \cup_\psi F'); \mathbb{Z}/2).$$

In the Pin^+ case the formula (4.3) shows that the composition along the bottom of the diagram is surjective, as we may always find a $g \in H^1(F, \partial F; \mathbb{Z}/2)$ that evaluates to 1 on $\sum a_i$.

In the Pin^- case, observe that given a tuple of elements in $\{[1], [3]\}^3$ we may write any element of $\mathbb{Z}/4$ as a sum of them using only coefficients 0 and 1. This observation along with the formula (4.4) shows that composition along the bottom of the diagram is surjective as long as F has genus ≥ 3 . $\square$

4.6. Homological stability. We will now use the above results to apply the main theorem of [18] in the case of non-orientable surfaces. The proofs necessarily use terminology introduced in [18], and we will not give these definitions again.

Theorem 4.19. *The moduli spaces of Pin^+ -surfaces exhibit homological stability. More precisely,*

- (i) Any $\alpha(n) : \mathcal{M}^{\text{Pin}^+}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^+}(S_{n+2,b-1})$ is a homology epimorphism in degrees $5* \leq n-1$ and a homology isomorphism in degrees $5* \leq n-6$.
- (ii) Any $\beta(n) : \mathcal{M}^{\text{Pin}^+}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^+}(S_{n,b+1})$ is a homology epimorphism in degrees $5* \leq n-2$, and a homology isomorphism in degrees $5* \leq n-7$. If one of the created boundary conditions is trivial, it is a split homology monomorphism in all degrees.
- (iii) Any $\gamma(n) : \mathcal{M}^{\text{Pin}^+}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^+}(S_{n,b-1})$ for $b \geq 2$ is a homology isomorphism in degrees $5* \leq n-2$, and a split homology epimorphism in all degrees.
- (iv) Any $\mu(n) : \mathcal{M}^{\text{Pin}^+}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^+}(S_{n+1,b})$ is a homology epimorphism in degrees $4* \leq n-1$, and a homology isomorphism in degrees $4* \leq n-5$.

Proof. In order to apply Theorem 7.2 of [18], we must verify a collection of conditions: that Pin^+ -structures stabilise on π_0 at genus h , and that they are k -trivial, and furthermore that they stabilise on π_0 at genus h' for projective planes, and that they are k' -trivial for projective planes. We claim that this is indeed so with $(h, k, h', k') = (4, 4, 4, 3)$, so solving the recurrence relations of [18, §7.2] with this data gives the stated stability ranges.

The path components of the moduli spaces of Pin^+ -structures stabilise at genus 4 by Proposition 4.14 and Lemma 4.18. Proof of 4-triviality follows from that in the (2-)Spin case, as all the diagrams which need to be checked involve only orientable surfaces, on which Pin^+ -structures are in canonical bijection with Spin-structures. Checking 3-triviality for projective planes amounts to producing, for each collection of Pin^+ -surfaces as in Figure 4 (where the two Pin^+ -surfaces obtained by going in the two possible ways around the diagram are isomorphic), a Pin^+ -surface taking the place of the dotted arrow.




FIGURE 4. The given data to show 3-triviality for projective planes. The two Pin^+ -surfaces obtained by composition are isomorphic, and we must provide a genus two Pin^+ -surface from A to C such that for each triangle the two obtained Pin^+ -surfaces are isomorphic.

We choose S_Δ so that $A(S_\Delta \circ S_B) = A(S_\Delta) + A(S_B) = A(S_L)$. This is possible as we can realise either value of A with a genus 2 surface. It now follows that

$$A(S_T \circ S_\Delta) = A(S_T) + A(S_\Delta) = A(S_T) + A(S_L) + A(S_B) = A(S_R).$$

As the surfaces S_R , S_L , $S_\Delta \circ S_B$ and $S_T \circ S_\Delta$ all have genus 3, the invariant A is complete, and hence $S_\Delta \circ S_B \cong S_L$ and $S_T \circ S_\Delta \cong S_R$ as Pin^+ -surfaces as required. $\square$

Theorem 4.20. *The moduli spaces of Pin^- -surfaces exhibit homological stability. More precisely,*

- (i) *Any $\alpha(n) : \mathcal{M}^{\text{Pin}^-}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^-}(S_{n+2,b-1})$ is a homology epimorphism in degrees $5* \leq n-1$ and a homology isomorphism in degrees $5* \leq n-6$.*
- (ii) *Any $\beta(n) : \mathcal{M}^{\text{Pin}^-}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^-}(S_{n,b+1})$ is a homology epimorphism in degrees $5* \leq n-2$ and a homology isomorphism in degrees $5* \leq n-7$. If one of the created boundary conditions is trivial, it is a split homology monomorphism in all degrees.*
- (iii) *Any $\gamma(n) : \mathcal{M}^{\text{Pin}^-}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^-}(S_{n,b-1})$ for $b \geq 2$ is a homology isomorphism in degrees $5* \leq n-2$, and a split homology epimorphism in all degrees.*
- (iv) *Any $\mu(n) : \mathcal{M}^{\text{Pin}^-}(S_{n,b}) \rightarrow \mathcal{M}^{\text{Pin}^-}(S_{n+1,b})$ is a homology epimorphism in degrees $5* \leq n-1$, and a homology isomorphism in degrees $5* \leq n-6$.*

Proof. In order to apply Theorem 7.2 of [18], we must verify a collection of conditions: that Pin^- -structures stabilise on π_0 at genus h , and that they are k -trivial, and furthermore that they stabilise on π_0 at genus h' for projective planes, and that they are k' -trivial for projective planes. We claim that this is indeed so with $(h, k, h', k') = (4, 4, 4, 4)$, so solving the recurrence relations of [18, §7.2] with this data gives the stated stability ranges.

The verifications are precisely the same as for Pin^+ -surfaces, with the sole exception that we only get 4-triviality for projective planes, not 3-triviality, because we cannot necessarily realise every value of the invariant A on $S_{2,1+1}$, though we can on $S_{3,1+1}$. $\square$

4.7. Homological stability for closing the last boundary. The above theorems give homological stability theorems for almost all the stabilisation maps one might require, except for that which closes off the last boundary component. The paper [18] contains a general result for closing the last boundary condition, whose hypotheses are satisfied in this case.

Theorem 4.21. *Any $\gamma(n) : \mathcal{M}^{\text{Pin}^\pm}(S_{n,1}; \delta) \rightarrow \mathcal{M}^{\text{Pin}^\pm}(\Sigma_n)$ is a homology isomorphism in degrees $5* \leq n-2$, and a homology epimorphism in degrees $5* \leq n+3$.*

Proof. We will verify the conditions for closeability for the tangential structures $\text{Pin}^\pm$, and so assume the language of [18]. The natural map $B_0^{\text{Pin}^\pm} \rightarrow B_0^O$, from the space of $\text{Pin}^\pm$ -structures on D^2 to the space of trivial tangential structures on D^2 , induces a bijection on π_0 as the fibre of $\theta : B\text{Pin}^\pm(2) \rightarrow BO(2)$ is path-connected.

Hence $\pi_0(C_0^{\text{Pin}^\pm}(F)) \cong \pi_0(C_0^O(F))$ for any surface F , and we showed in [18, §10.3.2] that the trivial tangential structure is closeable for non-orientable surfaces, so $\text{Pin}^\pm$ -structures are also. The theorem now follows from [18, Theorem 10.6]. $\square$

5. APPLICATIONS OF HOMOLOGY STABILITY FOR $\text{Pin}^\pm$ -SURFACES

Let us denote by $\mathbf{MTPin}^\pm(2)$ the Thom spectrum of the virtual bundle $-\gamma_2^{\text{Pin}^\pm} \rightarrow B\text{Pin}^\pm(2)$. There is a natural comparison map

$$\alpha : \coprod_{n \geq 1} \mathcal{M}^{\text{Pin}^\pm}(S_{n,1}; \delta) \longrightarrow \Omega^\infty \mathbf{MTPin}^\pm(2)$$

where δ is a boundary condition which bounds a disc. The left hand side admits the stricture of a topological monoid, under the “pair of pants” product, and Galatius

and the author [9] have shown that this map is a group-completion. In particular, applying the group-completion theorem [16] we obtain a homology equivalence

$$(5.1) \quad \mathbb{Z} \times \mathcal{M}^{\text{Pin}^\pm}(S_\infty) \longrightarrow \Omega^\infty \mathbf{MTPin}^\pm(2).$$

In particular, the group $\pi_0(\Omega^\infty \mathbf{MTPin}^+(2))$ is in natural isomorphism with $\mathbb{Z} \times \mathbb{Z}/2$ and the group $\pi_0(\Omega^\infty \mathbf{MTPin}^-(2))$ is in natural isomorphism with $\mathbb{Z} \times \mathbb{Z}/4$. Let us denote by $\Omega_\bullet^\infty \mathbf{MTPin}^\pm(2)$ those path components corresponding to 0 on the $\mathbb{Z}$ factor. Combining this with the homology stability theorem proved in this paper, we establish the following corollary.

Corollary 5.1. *The space $\mathcal{M}^{\text{Pin}^\pm}(S_{n,b}; \delta)$ has the integral homology of the infinite loop space $\Omega_\bullet^\infty \mathbf{MTPin}^\pm(2)$ in degrees $5* \leq n - 6$.*

In fact, one can do slightly better in the Pin^+ case, but we leave that to the reader. Taking fundamental groups, we obtain the following computational corollary.

Corollary 5.2. *The abelianisation of $\Gamma^{\text{Pin}^+}(S_{n,b}; \xi)$ is $\mathbb{Z}/2$ for $n \geq 9$. The abelianisation of $\Gamma^{\text{Pin}^-}(S_{n,b}; \xi)$ is $(\mathbb{Z}/2)^3$ for $n \geq 11$.*

Proof. By the homological stability theorems for these groups, their abelianisation in this range coincides with the first homology of the spaces $\Omega_0^\infty \mathbf{MTPin}^+(2)$ and $\Omega_0^\infty \mathbf{MTPin}^-(2)$ respectively. By Hurewicz' theorem, this coincides with the first homotopy group of these spaces. These are computed in Appendix A. $\square$

5.1. Stable homology of the $\text{Pin}^\pm$ mapping class groups. Let us briefly discuss the homology of the moduli spaces $\mathcal{M}^{\text{Pin}^\pm}(S_\infty)$, which coincides with the homology of the stable mapping class group $\Gamma^{\text{Pin}^\pm}(S_\infty)$. This will require certain calculations in homotopy theory which we have included as Appendix A. By Proposition A.3 and [17, §5.1] there are equivalences of $\mathbb{Z}[\frac{1}{2}]$ -local spectra

$$(5.2) \quad \mathbf{MTPin}^\pm(2) \left[\frac{1}{2} \right] \simeq \mathbf{MTO}(2) \left[\frac{1}{2} \right] \simeq \Sigma^\infty BO(2)_+ \left[\frac{1}{2} \right] \simeq \Sigma^\infty \mathbb{HP}_+^\infty \left[\frac{1}{2} \right]$$

and so isomorphisms on $\mathbb{Z}[\frac{1}{2}]$ -homology

$$H_*(\mathcal{M}^{\text{Pin}^\pm}(S_\infty); \mathbb{Z}[\frac{1}{2}]) \cong H_*(\mathcal{M}(S_\infty); \mathbb{Z}[\frac{1}{2}]) \cong H_*(Q(\mathbb{HP}_+^\infty); \mathbb{Z}[\frac{1}{2}]).$$

In particular, the rational cohomology ring is

$$H^*(\mathcal{M}^{\text{Pin}^\pm}(S_\infty); \mathbb{Q}) \cong H^*(\mathcal{M}(S_\infty); \mathbb{Q}) \cong \mathbb{Q}[\zeta_1, \zeta_2, \dots]$$

where the classes ζ_i in degree $4i$ are the characteristic classes introduced by Wahl [21] for unoriented surface bundles. For a surface bundle $S \rightarrow E \xrightarrow{\pi} B$ they may be defined as the Becker–Gottlieb transfer of the i -th power of the first Pontrjagin class of the vertical tangent bundle, that is, $\zeta_i(E) := \text{trf}_\pi^*(p_1(T^v E)^i) \in H^{4i}(B; \mathbb{Z})$.

Ebert and the author [6] studied the divisibility of $\zeta_i \in H^{4i}(\mathcal{M}(S_\infty); \mathbb{Z})$, and found them to be indivisible. By the first equivalence of (5.2) it is then clear that the classes $\zeta_i \in H^{4i}(\mathcal{M}^{\text{Pin}^\pm}(S_\infty); \mathbb{Z})$ are divisible at most by a power of 2.

Proposition 5.3. *The class $\zeta_i \in H^{4i}(\mathcal{M}^{\text{Pin}^-}(S_\infty); \mathbb{Z})$ is divisible by precisely 4^i . The class $\zeta_i \in H^{4i}(\mathcal{M}^{\text{Pin}^+}(S_\infty); \mathbb{Z})$ is divisible by 4^i , and by at most $2 \cdot 4^i$.*

Proof. Note that the class $p_1 \in H^*(B\text{Pin}^\pm(2); \mathbb{Z})$ is (uniquely) divisible by 4. Thus $\zeta_i = \text{trf}^*(p_1(T^v)^i) \in H^{4i}(\mathcal{M}^{\text{Pin}^\pm}(S_\infty); \mathbb{Z})$ is divisible by 4^i .

To see that it is not divisible further for Pin^- -structures, consider the projectivised vector bundle $\mathbb{RP}^2 \rightarrow \mathbb{P}(\gamma_3^{\text{Spin}}) \xrightarrow{\pi} B\text{Spin}(3)$. There is a vector bundle isomorphism $\pi^* \gamma_3^{\text{Spin}} \cong T^v \mathbb{P}(\gamma_3^{\text{Spin}}) \oplus L$ where L is the real line bundle characterised by the fact that it restricts to the tautological bundle on each fibre. By the

Leray–Hirsch theorem, the $\mathbb{F}_2$ -cohomology of the total space of this fibration is the free module

$$H^*(\mathbb{P}(\gamma_3^{\text{Spin}}); \mathbb{F}_2) \cong H^*(B\text{Spin}(3); \mathbb{F}_2)\langle 1, x, x^2 \rangle$$

where the class $x = w_1(L)$ satisfies $x^3 = 0$, and there is a similar decomposition for integral cohomology. The total Stiefel–Whitney class of the vertical tangent bundle is

$$w(T^v) = \pi^* w(\gamma_3^{\text{Spin}}) \cdot w(L)^{-1} = (1 + x + x^2),$$

so $w_1(T^v) = x$ and $w_2(T^v) = x^2$, and hence T^v admits a Pin^- -structure. Thus this bundle is formally classified by a map $B\text{Spin}(3) \rightarrow \mathcal{M}^{\text{Pin}^-}(S_1) \rightarrow \Omega^\infty \mathbf{MTPin}^-(2)$.

Recall that $H^*(B\text{Spin}(3); \mathbb{Z}) = \mathbb{Z}[\eta_1]$ where $4\eta_1 = p_1$. This may be seen as $\text{Spin}(3) = SU(2)$ so the cohomology ring is polynomial on a degree 4 generator $\eta_1 := -c_2(\gamma_2^{SU(2)})$, and $p_1(\gamma_3^{\text{Spin}}) = -4c_2 = 4\eta_1$. The calculation of the first Pontrjagin class follows by taking Chern classes for the identity $\gamma_3^{\text{Spin}} \otimes_{\mathbb{R}} \mathbb{C} = \text{Sym}^2(\gamma_2^{SU(2)})$, which we learnt from [5, Proposition 5.2.5]. Writing β for the Bockstein operation, we may compute¹ $\pi^*(p_1) = p_1(T^v \oplus L) = p_1(T^v) + \beta(w_1(T^v)) \cdot \beta(w_1(L))$ so that $p_1(T^v) = \pi^*(p_1) + \beta(x)^2$. The integral class $\beta(x)^2 \in H^4(\mathbb{P}(\gamma_3^{\text{Spin}}); \mathbb{Z}) = \mathbb{Z}\langle \eta_1 \cdot 1 \rangle$ is 2-torsion and hence zero. Thus $p_1(T^v) = \pi^*(p_1)$ so

$$\zeta_i = \text{trf}_\pi^*(p_1(T^v)^i) = \text{trf}_\pi^*(\pi^*(p_1^i)) = \chi(\mathbb{RP}^2) \cdot p_1^i = p_1^i = 4^i \eta_1^i \in H^{4i}(B\text{Spin}(3); \mathbb{Z})$$

which is divisible by precisely 4^i . Thus ζ_i is divisible by at most 4^i .

To pass from this information about the divisibility for bundles of $\mathbb{RP}^2$'s to divisibility in general, it is enough to note that the ζ_i are defined universally on $\Omega^\infty \mathbf{MTPin}^-(2)$, so by the above example must also be divisible by at most 4^i here. The homology equivalence (5.1) then implies the same divisibility in the cohomology of $\mathcal{M}^{\text{Pin}^-}(S_\infty)$.

To see that for Pin^+ -structures the class ζ_i is divisible by at most $2 \cdot 4^i$, note that pulling back ζ_i via $\Omega^\infty \mathbf{MTSpin}(2) \rightarrow \Omega^\infty \mathbf{MTPin}^+(2)$ gives the class κ_{2i} , which Ebert [5] has shown to be divisible by precisely 2^{2i+1} . $\square$

As only closed non-orientable surfaces of *even* Euler characteristic admit Pin^+ -structures, we cannot hope to find such a structure on the projectivisation of a rank 3 vector bundle, so making precise the upper bound on the divisibility of ζ_i in this case is more difficult.

APPENDIX A. COMPUTING $\pi_1(\mathbf{MTPin}^\pm(2))$

We first state the structure of the $\mathbb{F}_2$ -cohomology algebras of $B\text{Pin}^\pm(2)$, as algebras over the mod 2 Steenrod algebra $\mathcal{A}_2$. These may be computed from the Serre spectral sequence for the principal fibrations (4.1). In the Pin^- case one must consult the Eilenberg–Moore spectral sequence for the fibration $B\text{Pin}^-(2) \rightarrow BO(2) \rightarrow K(\mathbb{Z}/2, 2)$ to obtain the $\mathcal{A}_2$ -module structure.

Proposition A.1. *There is an isomorphism of rings*

$$H^*(B\text{Pin}^+(2); \mathbb{F}_2) \cong \mathbb{F}_2[w_1, x_2].$$

Here, w_1 is the first Stiefel–Whitney class of the universal Pin^+ -bundle, and the higher Stiefel–Whitney classes vanish. The $\mathcal{A}_2$ -module structure is determined by the ring structure and $Sq^1(x_2) = w_1 \cdot x_2$.

¹This is a general formula for the first Pontrjagin class of a rank 3 bundle which splits off a rank 1 subbundle: $p_1(\gamma_2 \oplus \gamma_1) = p_1(\gamma_2) + \beta(w_1(\gamma_2)) \cdot \beta(w_1(\gamma_1))$, which may be easily derived from the definition of Pontrjagin classes.

Proposition A.2. *There is an isomorphism of rings*

$$H^*(B\text{Pin}^-(2); \mathbb{F}_2) \cong \mathbb{F}_2[w_1, x_4]/(w_1^3).$$

Here, w_1 is the first Stiefel–Whitney class of the universal Pin^- -bundle, $w_2(\gamma_2^{\text{Pin}^-}) = w_1^2$, and the higher Stiefel–Whitney classes vanish. The class x_4 in degree 4 is the reduction of an integral class, and hence Sq^1 vanishes on it. Furthermore, Sq^2 also vanishes on it. This determines the structure as an $\mathcal{A}_2$ -module.

Proposition A.3. *There is an isomorphism $H^*(B\text{Pin}^\pm(2); \mathbb{Z}[\frac{1}{2}]) \cong \mathbb{Z}[\frac{1}{2}][p_1]$. The class p_1 in degree 4 is the first Pontrjagin class of the universal $\text{Pin}^\pm$ -bundle.*

Recall that $\mathbf{MTPin}^\pm(2)$ is defined to be the Thom spectrum of the virtual bundle $-\gamma_2^{\text{Pin}^\pm} \rightarrow B\text{Pin}^\pm(2)$. This has a Thom isomorphism in $\mathbb{F}_2$ -cohomology, so armed with the above data, one may easily compute the $\mathbb{F}_2$ -cohomology of $\mathbf{MTPin}^\pm(2)$ as an $\mathcal{A}_2$ -module, and hence compute the E_2 -page of the Adams spectral sequence in small degrees, which we include as Figure 5. Note also that by (5.2) the group $\pi_1(\mathbf{MTPin}^\pm(2))$ has only even torsion.




FIGURE 5. Partial E_2 -pages of the Adams spectral sequences converging to the 2-primary homotopy groups of the spectra $\mathbf{MTPin}^+(2)$ and $\mathbf{MTPin}^-(2)$ respectively. Vertical lines correspond to multiplication by $2 \in \pi_0(\mathbf{S})$, lines of slope 1 correspond to multiplication by $\eta \in \pi_1(\mathbf{S})$, and lines of slope $1/3$ correspond to multiplication by $\nu \in \pi_3(\mathbf{S})$. The diagram is complete to the left of the dotted line.

Theorem A.4. $\pi_1(\mathbf{MTPin}^+(2)) = \mathbb{Z}/2$.

Proof. The only possible differential landing in total degree 1 can be seen not to exist through the $\pi_*(\mathbf{S})$ -module structure. $\square$

Consider the cofibration sequence (see e.g. [8, Proposition 3.1]) of spectra

$$\mathbf{MTSpin}(2) \longrightarrow \mathbf{MTPin}^-(2) \longrightarrow \text{Th}(-\gamma_2^{\text{Pin}^-} \oplus \gamma_1^{\pm 1} \rightarrow B\text{Pin}^-(2)) =: \mathbf{C}$$

where $\gamma_1^{\pm 1}$ denotes the unique non-trivial real line bundle over $B\text{Pin}^-(2)$. The total Stiefel–Whitney class of $-\gamma_2^{\text{Pin}^-} \oplus \gamma_1^{\pm 1}$ is $(1 + w_1 + w_1^2)(1 + w_1) = 1$, which means that $H^*(\mathbf{C}; \mathbb{F}_2) \cong \Sigma^{-1}H^*(B\text{Pin}^-(2)_+; \mathbb{F}_2)$ as modules over the Steenrod algebra. As x_4 supports no non-trivial Steenrod operations, this module splits as

$$\Sigma^{-1}M \oplus \Sigma^3M \oplus \Sigma^7M \oplus \dots$$

where $M = \mathbb{F}_2\langle 1, w_1, w_1^2 \rangle$ is a module over the Steenrod algebra with unique operation $Sq^1(w_1) = w_1^2$.

Similarly, the total Stiefel–Whitney class of $-\gamma_2^{\text{Spin}} \rightarrow B\text{Spin}(2)$ is 1, so there is an isomorphism $H^*(\mathbf{MTSpin}(2); \mathbb{F}_2) \cong \Sigma^{-2}H^*(B\text{Spin}(2)_+; \mathbb{F}_2)$ of modules over the Steenrod algebra. This allows us to calculate the E_2 -pages of the Adams spectral sequences converging to the 2-primary homotopy groups of these spectra, which we include as Figure 6.




FIGURE 6. Partial E_2 -pages of the Adams spectral sequences converging to the 2-primary homotopy groups of the spectra $\mathbf{MTSpin}(2)$ and $\mathbf{C}$ respectively. The diagram is complete to the left of the dotted line.

Theorem A.5. $\pi_1(\mathbf{MTPin}^-(2)) = (\mathbb{Z}/2)^3$.

Proof. On the Adams E_2 -page in total degree 1, there is a $\mathbb{F}_2^2$ in filtration 1 and a $\mathbb{F}_2$ in filtration 2, and there are *no additive extensions*. In total degree 2 there is an $\mathbb{F}_2$ in filtration 0, which could potentially kill the element in total degree 1 filtration 2. We claim there is no such differential.

Consider the long exact sequence on homotopy coming from the cofibration sequence

$$\mathbf{MTSpin}(2) \longrightarrow \mathbf{MTPin}^-(2) \longrightarrow \mathbf{C},$$

$$\begin{aligned} \cdots \longrightarrow \pi_2(\mathbf{C}) \longrightarrow \pi_1(\mathbf{MTSpin}(2)) \xrightarrow{\neq 0} \pi_1(\mathbf{MTPin}^-(2)) \longrightarrow (\mathbb{Z}/2)^2 \xrightarrow{0} \\ \mathbb{Z} \oplus \mathbb{Z}/2 \longrightarrow \mathbb{Z} \oplus \mathbb{Z}/4 \longrightarrow (\mathbb{Z}/2)^2 \xrightarrow{0} \mathbb{Z}/2 \xrightarrow{\cong} \mathbb{Z}/2 \xrightarrow{0} \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \longrightarrow \mathbb{Z}/2. \end{aligned}$$

Working backwards from the end, the boundary map

$$\pi_{-1}(\mathbf{C}) = \mathbb{Z} \longrightarrow \pi_{-2}(\mathbf{MTSpin}(2)) = \mathbb{Z}$$

is multiplication by 2, as $\pi_{-2}(\mathbf{MTPin}^-(2)) = \mathbb{Z}/2$, and hence the previous map must be zero. Thus $\pi_{-1}(\mathbf{MTSpin}(2)) \rightarrow \pi_{-1}(\mathbf{MTPin}^-(2))$ is surjective and hence an isomorphism. The map

$$\pi_0(\mathbf{MTSpin}(2)) = \mathbb{Z} \oplus \mathbb{Z}/2 \longrightarrow \pi_0(\mathbf{MTPin}^-(2)) = \mathbb{Z} \oplus \mathbb{Z}/4$$

has cokernel $(\mathbb{Z}/2)^2$, so must be injective, and hence the previous map is zero. Note $\pi_1(\mathbf{MTSpin}(2))$ is generated by ν times a generator of $\pi_{-2}(\mathbf{MTSpin}(2))$, so it has non-trivial image in $\pi_1(\mathbf{MTPin}^-(2))$, and the result follows. $\square$

REFERENCES

- [1] Tilman Bauer. An infinite loop space structure on the nerve of spin bordism categories. *Q J Math.*, 55(2):117–133, 2004.
- [2] Ralph Cohen and Ib Madsen. Surfaces in a background space and the homology of mapping class group. *Proc. Symp. Pure Math.*, 80(1):43–76, 2009.
- [3] Ralph Cohen and Ib Madsen. Stability for closed surfaces in a background space. arXiv:1002.2498, 2010.

- [4] Alexander Degtyarev and Sergei Finashin. Pin-structures on surfaces and quadratic forms. *Turkish J. Math.*, 21(2):187–193, 1997.
- [5] Johannes Ebert. *Characteristic classes of spin surface bundles: Applications of the Madsen-Weiss theory*. PhD thesis, Bonner Mathematische Schriften 381, 2006.
- [6] Johannes Ebert and Oscar Randal-Williams. On the divisibility of characteristic classes of non-oriented surface bundles. *Topology Appl.*, 156(2):246–250, 2008.
- [7] Søren Galatius. Mod 2 homology of the stable spin mapping class group. *Math. Ann.*, 334:439 – 455, 2006.
- [8] Søren Galatius, Ib Madsen, Ulrike Tillmann, and Michael Weiss. The homotopy type of the cobordism category. *Acta Math.*, 202(2):195–239, 2009.
- [9] Søren Galatius and Oscar Randal-Williams. Monoids of moduli spaces of manifolds. *Geom. Topol.*, 14(3):1243–1302, 2010.
- [10] John L. Harer. Stability of the homology of the mapping class groups of orientable surfaces. *Ann. of Math. (2)*, 121(2):215–249, 1985.
- [11] John L. Harer. Stability of the homology of the moduli spaces of Riemann surfaces with spin structure. *Math. Ann.*, 287(2):323–334, 1990.
- [12] Dennis Johnson. Spin structures and quadratic forms on surfaces. *J. London Math. Soc. (2)*, 22(2):365–373, 1980.
- [13] W. B. R. Lickorish. Homeomorphisms of non-orientable two-manifolds. *Proc. Cambridge Philos. Soc.*, 59:307–317, 1963.
- [14] Ib Madsen and Michael Weiss. The stable moduli space of Riemann surfaces: Mumford’s conjecture. *Ann. of Math. (2)*, 165(3):843–941, 2007.
- [15] Gregor Masbaum. On representations of Spin mapping class groups arising in Spin TQFT. In *Geometry and physics (Aarhus, 1995)*, volume 184 of *Lecture Notes in Pure and Appl. Math.*, pages 197–207. Dekker, New York, 1997.
- [16] Dusa McDuff and Graeme Segal. Homology fibrations and the “group-completion” theorem. *Invent. Math.*, 31(3):279–284, 1975/76.
- [17] Oscar Randal-Williams. The homology of the stable nonorientable mapping class group. *Algebr. Geom. Topol.*, 8(3):1811–1832, 2008.
- [18] Oscar Randal-Williams. Resolutions of moduli spaces. arXiv:0909.4278, 2009.
- [19] Oscar Randal-Williams. Picard group of the moduli space of r -Spin Riemann surfaces. arXiv:1102.0715, 2011.
- [20] Roland Trapp. A linear representation of the mapping class group $\mathcal{M}$ and the theory of winding numbers. *Topology Appl.*, 43(1):47–64, 1992.
- [21] Nathalie Wahl. Homological stability for the mapping class groups of non-orientable surfaces. *Invent. Math.*, 171:389–424, 2008.

E-mail address: o.randal-williams@math.ku.dk

INSTITUT FOR MATEMATISKE FAG, UNIVERSITETSPARKEN 5, DK-2100 KØBENHAVN Ø, DENMARK